

Smooth Manifolds

Muchang Bahng

January 15, 2024

Contents

1	Smooth Manifolds	3
1.1	Topological Manifolds and Topological Properties	3
1.2	Smooth Structures	5
1.2.1	Local Coordinate Representations	9
1.2.2	Construction of a Smooth Manifold	9
1.3	Manifolds with Boundaries	10
2	Smooth Maps	11
2.1	Clarification of Multiple Meanings of Smoothness	14
2.2	Diffeomorphisms	14
2.2.1	Nondiffeomorphic Smooth Structures on Manifolds	15
2.3	Lie Groups	16
2.4	Smooth Covering Maps, Proper Maps	17
2.5	Partitions of Unity	17
2.5.1	Paracompactness	19
3	Tangent Vectors	19
3.1	Geometric Tangent Vectors	19
3.1.1	Tangent Vectors on Manifolds	21
3.2	Pushforwards	21
3.3	Computations in Coordinates	22
3.3.1	Change of Coordinates	24
3.4	Tangent Vectors to Curves	24
4	Vector Fields	25
4.1	The Tangent Bundle	25
4.2	Vector Fields on Manifolds	26
4.2.1	Pushforwards of Vector Fields	27
4.2.2	Vector Fields on Manifolds with Boundary	28
4.3	Lie Brackets	28
4.4	The Lie Algebra of a Lie Group	29
4.4.1	Induced Lie Algebra Homomorphisms	31
5	Vector (Fiber) Bundles	31
5.1	Categories and Functors	33
6	The Cotangent Bundle	36
6.1	Tangent Covectors on Manifolds	36
6.1.1	Basis Transformations	36
6.2	The Cotangent Bundle	37
6.3	The Differential of a Function	38
6.4	Pullbacks	39
6.5	Line Integrals	40
6.6	Conservative Covector Fields	42

7	Submersions, Immersions, and Embeddings	43
7.1	Maps of Constant Rank	43
7.2	The Inverse Function Theorem	43
8	Tensors	43
8.1	The Algebra of Tensors	44
8.2	Tensors and Tensor Fields on Manifolds	45
8.2.1	Pullbacks	46
8.3	Symmetric Tensors	46
8.4	Riemannian Metrics	47
8.4.1	Pseudo-Riemannian Metrics	47
9	Differential Forms	48
9.1	The Algebra of Alternating Tensors	48
9.2	The Wedge Product	49
9.3	Differential Forms on Manifolds	50
9.4	Exterior Derivatives	51
10	Orientations	52
10.1	Orientations on Vector Spaces	52
10.2	Orientations on Manifolds	52


Manifolds are invaluable in generalizing multiple ideas. For example, it allows us to generalize the concepts to calculus to spaces that are not only globally, but *locally* Euclidean.

1 Smooth Manifolds

1.1 Topological Manifolds and Topological Properties

Definition 1.1 (Topological Manifold, Coordinate Chart, Atlas). Suppose M is a topological space. M is a (real) *topological manifold of dimension N* if

1. M is Hausdorff: for every pair of points $p, q \in M$, there exists disjoint open subsets $U, V \subset M$ such that $p \in U, q \in V$.
2. M is second countable: There exists a countable basis for the topology of M .
3. M is locally homeomorphic to \mathbb{R}^N : There exists a covering of open sets in M where each open set is homeomorphic to an open set in \mathbb{R}^N . The visual below shows a 1-dimensional and 2-dimensional manifold.



Given open cover U_1, \dots, U_n with their respective homeomorphism maps $\varphi_1, \dots, \varphi_n$ where

$$\varphi_i : U_i \longrightarrow \mathbb{R}^n$$

and component maps

$$\varphi_i \equiv (x_{i1}, x_{i2}, \dots, x_{in})$$

their pairs (U_i, φ_i) are called *coordinate charts*. The collection of all coordinate charts

$$\mathcal{A} = \{(U_i, \varphi_i)\}_{i=1}^n$$

is called the *atlas* of M .

Note that the Hausdorff and second-countability condition is sometimes omitted from the definition of a manifold, but we keep it since:


1. many manifolds in nature have these properties.
2. we can deduce much more interesting properties about manifolds with these assumptions.

Theorem 1.1 (Product Manifolds). *Let M_1, M_2, \dots, M_n be topological manifolds of dimensions m_1, m_2, \dots, m_n , respectively. Then, the product space*

$$\prod_{i=1}^n M_i$$

is also a topological manifold with

$$\dim \prod_{i=1}^n M_i = \sum_{i=1}^n m_i$$




Example 1.1 (Graphs of Continuous Functions). Given a continuous function $f : U \rightarrow \mathbb{R}^m$, its graph

$$\Gamma(f) \equiv \{(x, y) \in \mathbb{R}^n \times \mathbb{R}^m \mid x \in U, y = f(x)\}$$

with the subspace topology is a topological manifold. Actually, this manifold is globally homeomorphic to an open set in \mathbb{R}^n , meaning that this is trivially a manifold.

Example 1.2 (Unit Sphere). The unit sphere S^n is a topological manifold since we can construct an atlas of charts formed by the stereographic projection. We show the stereographic projection for a 1-sphere.



Example 1.3. The n -sphere with the induced open ball topology of \mathbb{R}^{n+1} is not homeomorphic to \mathbb{R}^n since S^n is compact and \mathbb{R}^n is not. However, the n -sphere with one point $p \in S^n$ removed, $(S^n \setminus \{p\}, \tau_{\mathbb{R}^{n+1}}|_{S^n \setminus \{p\}})$ is homeomorphic to \mathbb{R}^n and is also not compact. We can visualize this homeomorphism by imagining the "hole" getting larger and stretching S^2 out to "look like" \mathbb{R}^2 .


Example 1.4. The n -torus, defined

$$\mathbb{T}^n \equiv \prod_n S^1$$

is a product manifold.

We end this section by stating a theorem that gives topological manifolds a nice basis structure. Recall that a subset U in a topological space X is said to be *precompact* if the closure \bar{U} is compact.

Lemma 1.2 (Basis of Topological Manifolds). *Every topological manifold has a countable basis of precompact coordinate balls.*







Lemma 1.3 (Fundamental Groups of Topological Manifolds). *The fundamental group of any topological manifold is countable.*

1.2 Smooth Structures

Note that even though topological manifolds bear some resemblance to Euclidean space, we cannot do calculus on them since derivatives are not invariant under homeomorphisms. For example, compare the two homeomorphisms $\varphi_1, \varphi_2 : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ (shown by visualizing the images of the grid lines), where

$$\varphi_1(u, v) = (u, v)$$

$$\varphi_2(u, v) = (u^{1/3}, v^{1/3})$$



Given function $f : M \rightarrow \mathbb{R}^m$, if topological manifold M has chart φ_1 , f will be considered smooth at $\varphi_1^{-1}(0)$, but if M has chart φ_2 , f will not be considered smooth at $\varphi_2^{-1}(0)$.

Clearly, it is a problem if two different charts around a point gives contradicting things about its smoothness. We can fix this by introducing a structure that makes smoothness invariant. First, recall the definition of a diffeomorphism.


Definition 1.2 (Diffeomorphism). If smooth $f : U \subset \mathbb{R}^n \rightarrow V \subset \mathbb{R}^m$ is bijective and has a smooth inverse map, then f is said to be a *diffeomorphism*. That is, a diffeomorphism is a homeomorphism where both the function and its inverse is smooth. More specifically,

1. If f, f^{-1} is of class C^k , then it is a C^k -diffeomorphism.
2. f, f^{-1} is of class C^∞ , then it is a C^∞ -diffeomorphism, or a smooth diffeomorphism.

Definition 1.3 (Transition Maps, Smooth Atlases). Let M be a topological manifold with $(U, \varphi), (V, \psi)$ two charts such that $U \cap V \neq \emptyset$. Then, the composite map (between Euclidean spaces)

$$\psi \circ \varphi^{-1} : \varphi(U \cap V) \rightarrow \psi(U \cap V)$$

is called the *transition map* from φ to ψ .




The two charts are said to be smoothly (or C^k) compatible if either

1. $U \cap V = \emptyset$, or
2. the transition map $\psi \circ \varphi^{-1}$ is a smooth (or C^k) diffeomorphism. Since $\psi \circ \varphi^{-1}$ is a map between Euclidean spaces, this means that $\psi \circ \varphi^{-1}$ has continuous partial derivatives.

An atlas \mathcal{A} is called a *smooth (C^k) atlas* if any 2 charts in \mathcal{A} are smoothly (C^k) compatible with each other.

Example 1.5. A transition map can be defined between two stereographic projections of the 1-sphere $\delta_1 : S^1 \setminus \{p\} \rightarrow \mathbb{R}$ and $\delta_2 : S^1 \setminus \{q\} \rightarrow \mathbb{R}$, where p and q are diametrically opposite points of S^1 . By placing \mathbb{R} within S^1 orthogonal to the line segment PQ and intersecting the center of S^1 , we find that the transition function $\delta_1 \circ \delta_2^{-1}(n) = \delta_2 \circ \delta_1^{-1}(n) = \frac{1}{n}$.




Let's go back to our example problem earlier. Given an open neighborhood M around $0 \in \mathbb{R}^2$, the two (global) charts φ_1, φ_2 are not smoothly compatible since

$$(\varphi_2 \circ \varphi_1^{-1})(u, v) = (u^{1/3}, v^{1/3})$$

is not smooth. Therefore, they cannot be a part of the same smooth atlas.

Our previous problem hints at another one: Which atlas is the "right" one? In general, there will be many possible choices of atlases that give the "same" smooth structure. The 3 following atlases shown below (consisting of one global chart, for simplicity), are all viable.



In the visual above, notice that

1. $\mathcal{A}_1, \mathcal{A}_2, \mathcal{A}_3$ are all smooth atlases (trivially).
2. φ_1 and φ_2 are smoothly compatible $\implies \mathcal{A}_{12} = \{(M, \varphi_1), (M, \varphi_2)\}$ is a smooth atlas.
3. φ_3 is not smoothly compatible with either φ_1 nor φ_2 . This implies that $\mathcal{A}_{13} = \{(M, \varphi_1), (M, \varphi_3)\}, \mathcal{A}_{23} = \{(M, \varphi_2), (M, \varphi_3)\}$ are not smooth atlases.

Clearly, φ_1 and φ_2 are closely related as they are smoothly compatible. Furthermore,

$$\mathcal{A}_1 \subset \mathcal{A}_{12}, \quad \mathcal{A}_2 \subset \mathcal{A}_{12}$$

Now, imagine an atlas that contains $(M, \varphi_1), (M, \varphi_2)$, and all other possible charts that are smoothly compatible with φ_1 and φ_2 . This creates a "maximal" smooth atlas.


Definition 1.4 (Maximal Smooth Atlas). A *maximal smooth* (C^k) *atlas* is a smooth (C^k) atlas that is not contained in any strictly larger smooth atlas. In other words, it is the largest possible atlas in which every chart is smoothly (C^k) compatible with one another.

The existence of this maximal smooth (C^k) atlas wouldn't be as helpful without the following lemma, which allows us to easily describe it.

Lemma 1.4 (Induced Maximal Smooth Atlases of Topological Manifolds). *Let M be a topological n -manifold.*

1. *Every smooth atlas for M is contained in a unique maximal smooth atlas.*
2. *Two smooth atlases for M determine the same maximal smooth atlas if and only if their union is a smooth atlas.*

That is, let $A_1, \dots, A_a, B_1, \dots, B_b, C_1, \dots, C_c$ be a collection of smooth atlases on M such that any union of the A_i 's, any union of the B_i 's, and any union of the C_i 's, are smooth atlases. Then, we can imagine all of the A_i 's as subsets of the unique maximal atlas A^ , all of the B_i 's as subsets of the unique maximal atlas B^* , and all of the C_i 's as subsets of the unique maximal atlas C^* .*



Therefore, there can exist many smooth structures for topological manifold M .

From the lemma, we can clearly see that the atlases form an equivalence class, one for smooth structure.

Definition 1.5 (Smooth Equivalence Relation). Two atlases $\mathcal{A}_1, \mathcal{A}_2$ are *smoothly* (C^k) *equivalent* if $\mathcal{A}_1 \cup \mathcal{A}_2$ forms a smooth (C^k) atlas. Note that C^k -equivalence is a relation, and so the collection of all atlases that are C^k -equivalent forms a C^k -equivalence class, which is actually determined uniquely by the maximal atlas.

Definition 1.6 (Smooth (C^k) Structure, Smooth (C^k) Manifolds). Three closely related definitions:

1. A *smooth* (C^k) *structure* on a topological n -manifold M is a maximal smooth (C^k) atlas.
2. A *smooth* (C^k) *manifold* is a pair (M, \mathcal{A}) , M being a topological manifold and \mathcal{A} a smooth (C^k) structure.
3. A chart, or a coordinate map, of a smooth manifold, is called a *smooth chart*.

Three final things to mention:

1. There exist topological manifolds that admit no smooth structures at all. So, we cannot add a smooth structure to every topological manifold.
2. It is generally not convenient to define a smooth structure by explicitly describing a maximal smooth atlas, since such an atlas contains many charts. Fortunately, by the previous lemma, we only need to specify *some* smooth atlas, which will induce a maximal smooth atlas.

Specify $A_i \implies$ Smooth Structure = A^*

Specify $B_i \implies$ Smooth Structure = B^*

Specify $C_i \implies$ Smooth Structure = C^*

3. In many cases, one proves that a topological manifold M has a smooth atlas by directly computing and seeing that $\psi \circ \varphi^{-1}$ is smooth, for every pair ψ, φ in the atlas. Clearly, all of the $\psi \circ \varphi^{-1}$ are homeomorphisms (as compositions of homeomorphisms), and since we've proved them to be smooth, it automatically follows that they are diffeomorphisms. j

Definition 1.7 (Other Classes of Manifolds). Let M be a manifold.

1. A C^0 manifold is just a topological manifold.
2. If the transition mappings of M can be expressed as real analytic (i.e. expressible as a convergent power series in a neighborhood of each point), then M is said to have a C^ω structure, making M a *real-analytic manifold*.
3. If M has an even dimension $2m$, then we can use the fact that $\mathbb{R}^{2m} \simeq \mathbb{C}^m$ to define a C^m structure, making M a *complex manifold*.

Example 1.6. Consider \mathbb{R} with the charts (\mathbb{R}, id) and (\mathbb{R}, x^3) . Each of these charts cover \mathbb{R} , but they are not C^∞ -compatible (since $\sqrt[3]{x}$ is not C^∞), which means that they generate different maximal atlases. In fact, in \mathbb{R} , there are an infinitely many non-compatible maximal atlases each giving the topological space \mathbb{R} the structure of a differentiable manifold.

We can actually prove a stronger theorem.


Theorem 1.5 (Distinct Smooth Structures on Positive-Dimensional Smooth Manifolds). *Let M be a nonempty topological manifold of dimension $n \geq 1$. If M has a smooth structure, then it has uncountably many distinct ones.*

1.2.1 Local Coordinate Representations

We describe how one usually thinks about coordinate charts on a smooth manifold. Once we choose a smooth chart (U, φ) on M , the coordinate map

$$\varphi : U \longrightarrow \tilde{U} \subset \mathbb{R}^n$$

can be thought of as giving an *identification* between U and \tilde{U} . Using this identification, we can think of U simultaneously as an open subset of M and as an open subset of \mathbb{R}^n . You can visualize this identification by thinking of a "grid" drawn on U representing the inverse images of the coordinate lines under φ .



Using this identification, we can represent point $p \in U$ by its coordinates

$$(x^1, x^2, \dots, x^n) = \varphi(p)$$

and think of this n -tuple as *being* the point p (even though p , as an abstract point, really has no coordinates). This is typically expressed by saying that (x^1, \dots, x^n) is the (local) coordinate representation for p or " $p = (x^1, \dots, x^n)$ in local coordinates." Note that the curves (lines) do not necessarily have to be rectilinear. The projected lines from Euclidean space onto the manifold can be of any shape, including polar, as long as it is a homeomorphism.

Definition 1.8 (Einstein Summation Notation). Due to the abundance of summations in this chapter, we will abbreviate such a sum as such

$$\sum_i x^i E_i = x^i E_i$$

1.2.2 Construction of a Smooth Manifold

When defining a smooth manifold, we start with a topological space and check that it is a topological manifold, and then we specify a smooth structure. The following lemma combines these steps into one.

Lemma 1.6 (Smooth Manifold Construction Lemma). *Let M be a set, and suppose we are given a collection $\{U_\alpha\}$ of subsets of M , together with an injective map $\varphi_\alpha : U_\alpha \longrightarrow \mathbb{R}^n$ for each α , such that the following properties are satisfied:*

1. For each α , $\varphi_\alpha(U_\alpha)$ is an open subset of \mathbb{R}^n .
2. For each α and β , $\varphi_\alpha(U_\alpha \cap U_\beta)$ and $\varphi_\beta(U_\alpha \cap U_\beta)$ are open in \mathbb{R}^n .

3. Whenever $U_\alpha \cap U_\beta \neq \emptyset$, $\varphi_\alpha \circ \varphi_\beta^{-1} : \varphi_\beta(U_\alpha \cap U_\beta) \rightarrow \varphi_\alpha(U_\alpha \cap U_\beta)$ is a diffeomorphism.
4. M is second countable.
5. M is Hausdorff.

Then, M has a unique smooth manifold structure such that each $(U_\alpha, \varphi_\alpha)$ is a smooth chart.


1.3 Manifolds with Boundaries

We may come across manifolds which have a "boundary" of some sort, such as the closed unit ball in \mathbb{R}^n and the closed upper hemisphere in S^n .

Definition 1.9 (Euclidean Upper Half-Space). The closed n -dimensional upper half-space $\mathbb{H}^n \subset \mathbb{R}^n$ is defined

$$\mathbb{H}^n \equiv \{(x^1, x^2, \dots, x^n) \in \mathbb{R}^n \mid x^n \geq 0\}$$

\mathbb{H}^2 and \mathbb{H}^3 are shown below.




Clearly, \mathbb{H}^n has the subspace topology induced by that of \mathbb{R}^n , allowing charts to map open neighborhoods of boundary points. We also define

$$\begin{aligned} \text{Int}(\mathbb{H}^n) &\equiv \{(x^1, x^2, \dots, x^n) \in \mathbb{R}^n \mid x^n > 0\} \\ \partial(\mathbb{H}^n) &\equiv \{(x^1, x^2, \dots, x^n) \in \mathbb{R}^n \mid x^n = 0\} \end{aligned}$$

Definition 1.10 (Smooth Maps between Subsets). Note that a smooth map from an arbitrary subset $A \subset \mathbb{R}^n$ to \mathbb{R}^k is defined to be a map that admits a smooth extension to an open neighborhood of each point.

Definition 1.11 (Topological Manifold with Boundary). An n -dimensional topological manifold with boundary is a second-countable Hausdorff space M that is locally homeomorphic to \mathbb{H}^n . An open subset $U \subset M$ together with a homeomorphism φ from U to an open subset of \mathbb{H}^n is called a chart.

1. (U, φ) is an interior chart if $\varphi(U) \subset \text{Int}(\mathbb{H}^n)$.
2. (U, φ) is a boundary chart if $\varphi(U) \cap \partial\mathbb{H}^n \neq \emptyset$



A point $p \in M$ is a *boundary point* if its image under some smooth chart is in $\partial\mathbb{H}^n$; the set of all such boundary points is denoted ∂M . The set of all interior points of M is denoted $\text{Int}(M)$. Furthermore, it turns out that

$$M = \text{Int}(M) \sqcup \partial M$$

That is, M can be partitioned into its interior and boundary.

To define a smooth structure on a manifold with boundary, we must be able to define smooth maps that encompasses cases between boundary charts.

Definition 1.12. Thus, if U is an open subset of \mathbb{H}^n , a map $F : U \rightarrow \mathbb{R}^k$ is smooth if for each $x \in U$ there exists an open neighborhood $V \subset \mathbb{R}^n$ of x and a smooth map (extension) $\tilde{F} : V \rightarrow \mathbb{R}^k$ that agrees with F on $V \cap \mathbb{H}^n$.

Definition 1.13 (Smooth Manifold with Boundary). Let M be a topological manifold with boundary. A *smooth structure* for M is defined to be a maximal smooth atlas, i.e. a collection of charts whose domains cover M and whose transition maps (and inverses) are smooth as in it admits smooth extensions. With such a structure, M is called a *smooth manifold with boundary*.

Note that the boundary of manifold M and the boundary of M as a subset of a bigger topological space are two completely different sets.

Example 1.7. Let B^2 be the open unit disk in \mathbb{R}^2 . Then, \bar{B}^2 , the closed unit disk, is a smooth manifold with boundary, with the boundary being the circle S^1 . However, if we interpret B^2 as a topological subspace

1. \mathbb{R}^2 , the topological boundary is S^1 .
2. \mathbb{R}^3 , the topological boundary is \bar{B}^2 .
3. \bar{B}^2 , the topological boundary is \emptyset .

Furthermore, every smooth n -manifold can be considered a smooth n -manifold with boundary by composing each chart mapping with a diffeomorphism from \mathbb{R}^n to \mathbb{H}^n such as

$$(x^1, \dots, x^{n-1}, x^n) \mapsto (x^1, \dots, x^{n-1}, e^{x^n})$$


This modifies all manifold charts to take its values in $\text{Int}(\mathbb{H}^n)$ without affecting the smooth compatibility condition. Also, given a smooth n -manifold with boundary M , $\text{Int}(M)$ is also a topological n -manifold since the subfamily of all smooth interior charts is also a smooth atlas.

2 Smooth Maps

Definition 2.1 (Smooth Maps from Manifolds to Euclidean Space). If M is a smooth n -manifold, a function $f : M \rightarrow \mathbb{R}^k$ is said to be *smooth* if for every $p \in M$, there exists a smooth chart (U, φ) for M whose domain contains p and such that the composite function, called the *coordinate representation of f*

$$\hat{f} = f \circ \varphi^{-1} : \varphi(U) \subset \mathbb{R}^n \rightarrow \mathbb{R}^k$$

is smooth.



By definition, f is smooth if and only if its coordinate representation is smooth in some smooth chart around every point.

The set of smooth real-valued functions $f : M \rightarrow \mathbb{R}$ is denoted as $C^\infty(M)$. Since sums and constant multiples of smooth functions are smooth, $C^\infty(M)$ is a vector space.

Definition 2.2 (Smooth Maps between Manifolds). Let M, N be smooth m, n -manifolds, and let $F : M \rightarrow N$ be any map. Then, F is a *smooth map* if for every $p \in M$, there exist smooth charts (U, φ) containing p and (V, ψ) containing $F(p)$ such that $F(U) \subset V$ and the composite map

$$\psi \circ F \circ \varphi^{-1} : \varphi(U) \subset \mathbb{R}^m \rightarrow \psi(V) \subset \mathbb{R}^n$$


is smooth (in the regular Euclidean sense). The abstraction of F does not do us much good computation wise, so more often than not, we look at the *coordinate representation of F*

$$\hat{F} \equiv \psi \circ F \circ \varphi^{-1}$$

In here, \hat{F} is really just a representation of the abstract map F in specific local coordinates determined by φ and ψ . Once φ, ψ are determined, we can often ignore the distinction between F and \hat{F} .

Application-wise, there are three ways to prove that a particular map is smooth:

1. Write the map in smooth local coordinates and recognize its component functions as compositions of smooth elementary functions.
2. Exhibit the map as a composition of known smooth maps.
3. Use some special-purpose theorem that applies to the particular case.



Theorem 2.1 (Properties of Smooth Maps between Manifolds). *Given M, N smooth m, n -manifolds, with arbitrary map $F : M \rightarrow N$, we have the following properties:*

1. *If F is smooth, then F is continuous.*
2. *The composition of smooth maps between smooth manifolds is smooth.*
3. *If there exists an open cover $\{U_\alpha\}_\alpha$ of M and smooth maps $F_\alpha : U_\alpha \rightarrow N$ such that they agree on overlaps*

$$F_\alpha|_{U_\alpha \cap U_\beta} = F_\beta|_{U_\alpha \cap U_\beta} \text{ for all } \alpha, \beta$$

then there exists a unique smooth map $F : M \rightarrow N$ such that F agrees with all the F_α 's.

The last property is convenient for when we wish to construct a global smooth map from local ones.

Proof. We prove the first two properties:

1. Suppose $F : M \rightarrow N$ is smooth. The definition of smoothness guarantees that for every $p \in M$, we can choose smooth charts (U, φ) containing p and (V, ψ) containing $F(p)$ such that $F(U) \subset V$ and $\psi \circ F \circ \varphi^{-1} : \varphi(U) \rightarrow \psi(V)$ is smooth, hence continuous. Since φ and ψ are homeomorphisms, this implies that


$$F|_U = \psi^{-1} \circ (\psi \circ F \circ \varphi^{-1}) \circ \varphi : U \rightarrow V$$

is continuous, as a composition of continuous maps. Since F is continuous at each point, it is continuous on M .

2. Looking at the diagram below, we can see that since F, G are smooth, the mappings (between Euclidean spaces) $\theta \circ F \circ \varphi^{-1}$ and $\psi \circ G \circ \theta^{-1}$ are smooth, meaning that their composition

$$(\psi \circ G \circ \theta^{-1}) \circ (\theta \circ F \circ \varphi^{-1}) = \psi \circ (G \circ F) \circ \varphi^{-1}$$

is smooth. By definition, this means that $G \circ F$ is smooth.




■

2.1 Clarification of Multiple Meanings of Smoothness

We emphasize that the smoothness of a map F between manifolds depends **only** on the smoothness of its coordinate representation \hat{F} ! This warning is more clearly explained through the following example.

Example 2.1 (Mappings between Lines with Different Smooth Structures). *Let us have manifold \mathbb{R} with the standard smooth structure determined by global mapping $id: \mathbb{R} \rightarrow \mathbb{R}$. Let us have a second manifold $\tilde{\mathbb{R}}$ with smooth structure determined by global mapping $\psi(x) = x^{1/3}$. Now, let us have mappings (between manifolds) $F_1, F_2, F_3: \mathbb{R} \rightarrow \tilde{\mathbb{R}}$ where $F_1(x) = x, F_2(x) = x^3, F_3(x) = x^6$, as shown below.*



Then,

$$\begin{aligned} \psi \circ F_1 \circ id^{-1}(x) &= x^{1/3} \implies F_1 \text{ not smooth} \\ \psi \circ F_2 \circ id^{-1}(x) &= x \implies F_2 \text{ smooth} \\ \psi \circ F_3 \circ id^{-1}(x) &= x^2 \implies F_3 \text{ smooth} \end{aligned}$$

Note that the smoothness of the F_i 's when interpreting them as mappings between manifolds gives results compared to when we interpret them as regular mappings between Euclidean space. For example, F_1 is not smooth in the manifold sense even though it is clearly smooth in the Euclidean sense (since it is the identity mapping).

2.2 Diffeomorphisms

Definition 2.3 (Diffeomorphism). A *diffeomorphism* between smooth manifolds M and N is a smooth bijective map $F: M \rightarrow N$ that has a smooth inverse. It is said that M and N are *diffeomorphic* if there exists a diffeomorphism between them, denoted as

$$M \approx N$$

This is in fact an equivalence relation. Just as two topological spaces are considered to be the same if they are homeomorphic, two smooth manifolds are essentially indistinguishable if they are diffeomorphic. Clearly, diffeomorphisms preserve the dimensionalities of M and N .


Clearly, if M is any smooth manifold and (U, φ) is a smooth coordinate chart on M , then $\varphi: U \rightarrow \varphi(U) \subset \mathbb{R}^n$ is a diffeomorphism (by definition).

Example 2.2. *Let B^n be the open unit ball in \mathbb{R}^n (each with the standard smooth structure). Then, the map*

$$F: B^n \rightarrow \mathbb{R}^n, F(x) \equiv \frac{x}{1 - \|x\|^2}$$

is a diffeomorphism.

Example 2.3 (Mappings between Lines with Different Smooth Structures). *In the diagram below regarding a previous example, we have the three maps $F_1, F_2, F_3: \mathbb{R} \rightarrow \tilde{\mathbb{R}}$, where \mathbb{R} is a smooth manifold with the standard smooth structure, and $\tilde{\mathbb{R}}$ is a smooth manifold with the smooth structure consisting of ψ .*



To find out whether F_i is a diffeomorphism, we must confirm that F_i is bijective, smooth, and F_i^{-1} is smooth.

1. F_1 is clearly bijective. $\psi \circ F_1 \circ id^{-1}(x) = x^{1/3}$, so F_1 is not smooth. $id \circ F_1^{-1} \circ \psi^{-1} = x^3$, so F_1^{-1} is smooth.
2. F_2 is clearly bijective. $\psi \circ F_2 \circ id^{-1}(x) = x$, so F_2 is smooth. $id \circ F_2^{-1} \circ \psi^{-1} = x$, so F_2^{-1} is smooth.
3. F_3 is not bijective. $\psi \circ F_3 \circ id^{-1}(x) = x^2$, so F_3 is smooth. Since F_3 is not injective, F_3^{-1} is not well-defined.

Therefore, F_2 is the only diffeomorphism out of the three observed functions.

Definition 2.4 (Local Diffeomorphism). $F : M \rightarrow N$ is called a *local diffeomorphism* if every point $p \in M$ has a neighborhood U such that $F(U)$ is open in N and

$$F|_U : U \rightarrow F(U)$$

is a diffeomorphism.

2.2.1 Nondiffeomorphic Smooth Structures on Manifolds

We already found out that there are many distinct structures on a positive-dimensional smooth manifold (in fact, an uncountable number of them). Furthermore, we have seen in this section that there may exist a diffeomorphism between the two copies of the topological manifold, each with distinct smooth structures ($F_2(x) = x^3$ was a diffeomorphism between (\mathbb{R}, id) and (\mathbb{R}, ψ)).

This leads to the more interesting question of whether a given topological manifold admits smooth structures that are *not* diffeomorphic to each other (as in, there exists no diffeomorphism F between two copies of a given topological manifold, each with distinct smooth structures).

Definition 2.5 (Diffeomorphic Smooth Structures). Given a topological manifold M , let \mathcal{A}_1 and \mathcal{A}_2 any two smooth structures on M . Then,

$$(M, \mathcal{A}_1) \approx (M, \mathcal{A}_2)$$


if there exists some diffeomorphism $F : (M, \mathcal{A}_1) \rightarrow (M, \mathcal{A}_2)$. If the underlying manifold M is known, then we write this more concisely as

$$\mathcal{A}_1 \approx \mathcal{A}_2$$

and say that

1. \mathcal{A}_1 and \mathcal{A}_2 are diffeomorphic smooth structures on M , or
2. \mathcal{A}_1 and \mathcal{A}_2 are identical smooth structures on M up to diffeomorphism.

In fact, the properties of smooth mappings imply that this relation \approx endowed on the set of all smooth structures of M forms equivalence classes of diffeomorphic smooth structures. Visually, given that A^*, B^*, C^*, \dots are smooth structures on M , they can be classified further into their respective diffeomorphism classes.



Example 2.4 (Diffeomorphic Smooth Structures of \mathbb{R}). Since $F_2 = x^3$ is a diffeomorphism between (\mathbb{R}, id) and $(\tilde{\mathbb{R}}, \psi)$, the two atlases $\{\mathbb{R}, id\}$ and $\{\tilde{\mathbb{R}}, \psi\}$ are diffeomorphic smooth structures:

$$\{\mathbb{R}, id\} \approx \{\tilde{\mathbb{R}}, \psi\}$$

It turns out that every smooth structure of \mathbb{R} is equivalent in this sense. That is, there is only one smooth structure on \mathbb{R} up to diffeomorphism.

Theorem 2.2 (Classification of Nondiffeomorphic Smooth Structures on \mathbb{R}^n). The classification of smooth structures on Euclidean space is as follows:

1. When $n \neq 4$, \mathbb{R}^n has one unique smooth structure up to diffeomorphism.
2. \mathbb{R}^4 has uncountably many distinct smooth structures, no two of which are diffeomorphic to each other! The study of exotic \mathbb{R}^4 s arises from this phenomenon.

For compact manifolds, the situation is even more fascinating.

Theorem 2.3 (Classification of Nondiffeomorphic Smooth Structure on S^n). The table below details the number of unique smooth structures on S^n (for n up to 12) up to diffeomorphism.


n	1	2	3	4	5	6	7	8	9	10	11	12
Smooth Struc. on S^n	1	1	1	?	1	1	28	2	8	6	992	1

Notice that the number of smooth structures on the exotic 4-sphere is still unanswered.


2.3 Lie Groups

Definition 2.6 (Lie Group). A Lie group is a smooth manifold G that is also a group in the algebraic sense, with the property that

1. the multiplication map $m : G \times G \rightarrow G$, $m(g, h) \equiv gh$ is smooth. That is, the function $\phi \circ m \circ (\varphi \times \psi)^{-1} : \mathbb{R}^{2m} \rightarrow \mathbb{R}^m$ in the visual below is smooth in the Euclidean sense.



- the inversion map $i : G \rightarrow G$, $i(g) \equiv g^{-1}$ is smooth. That is, the function $\theta \circ i \circ \varphi^{-1} : \mathbb{R}^m \rightarrow \mathbb{R}^m$ in the visual below is smooth in the Euclidean sense.



Definition 2.7 (Lie Group Homomorphism). If G and H are Lie groups, a *Lie group homomorphism* from G to H is a smooth map $F : G \rightarrow H$ that is also a group homomorphism. The smoothness of F (a smooth property) and its preservice of the algebraic structure of G (an algebraic property) can be comprehensively shown in the commutative diagram below.

$$\begin{array}{ccccc}
 \mathbb{R}^m \times \mathbb{R}^m & \xleftarrow{\varphi \times \varphi} & G \times G & \xrightarrow{m_1} & G & \xrightarrow{\varphi} & \mathbb{R}^m \\
 & & \downarrow F \times F & & \downarrow F & & \\
 \mathbb{R}^n \times \mathbb{R}^n & \xleftarrow{\psi \times \psi} & H \times H & \xrightarrow{m_2} & H & \xrightarrow{\psi} & \mathbb{R}^n
 \end{array}$$

Note that

- $\varphi \times m_1 \times (\varphi \times \varphi)^{-1} : \mathbb{R}^m \times \mathbb{R}^m \rightarrow \mathbb{R}^m$ and $\psi \circ m_2 \circ (\psi \times \psi)^{-1} : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ are smooth since G and H are Lie groups.
- $F \circ m_1 : G \times G \rightarrow H$ and $m_2 \circ (F \times F) : G \times G \rightarrow H$ are smooth since F is a Lie group homomorphism. Note that since $F \circ m_1$ and $m_2 \circ (F \times F)$ are smooth maps between manifolds, it really means that the maps

$$\begin{aligned}
 &\psi \circ F \circ m_1 \circ (\varphi \times \varphi)^{-1} : \mathbb{R}^m \times \mathbb{R}^m \rightarrow \mathbb{R}^n \\
 &\psi \circ m_2 \circ (F \times F) \circ (\varphi \times \varphi)^{-1} : \mathbb{R}^m \times \mathbb{R}^m \rightarrow \mathbb{R}^n
 \end{aligned}$$

are smooth in the Euclidean sense.

F is called a *Lie group isomorphism* if it is also a diffeomorphism, which implies that it has an inverse that is also a Lie group homomorphism. This means that G and H are *isomorphic Lie groups*, meaning that they are indistinguishable as topological spaces, smooth manifolds, and as algebraic structures.

2.4 Smooth Covering Maps, Proper Maps

TBD TBD

2.5 Partitions of Unity

Recall the Gluing Lemma from topology, which allows us to construct continuous maps by "gluing together" maps defined on subspaces.

Lemma 2.4 (Pasting Lemma, Gluing Lemma). *Let $X = A \cup B$, where A, B are closed in X . Let $f : A \rightarrow Y$ and $g : B \rightarrow Y$ be continuous. If*

$$f(x) = g(x) \text{ for all } x \in A \cap B$$

Then f and g can be combined to form a continuous function $h : X \rightarrow Y$, defined

$$h(x) \equiv \begin{cases} f(x) & x \in A \setminus B \\ f(x) \text{ or } g(x) & x \in A \cap B \\ g(x) & x \in B \setminus A \end{cases}$$

For smooth manifolds, however, the gluing lemma is of limited usefulness, since the produced map, while continuous, is rarely smooth. Observe the following example.

Example 2.5. *Given two functions $f_+ : [0, \infty) \rightarrow \mathbb{R}$ and $f_- : (-\infty, 0] \rightarrow \mathbb{R}$ defined by*

$$\begin{aligned} f_+(x) &= +x, x \in [0, \infty) \\ f_-(x) &= -x, x \in (-\infty, 0] \end{aligned}$$

are both smooth and agree at the point 0 where they overlap, but the continuous map $f : \mathbb{R} \rightarrow \mathbb{R}$ that they define, $f(x) = |x|$, is not smooth at the origin.

Partitions of unity solves this problem as tools for patching together local smooth maps into global ones. We must first establish the existence of smooth functions that are positive in a specified part of a manifold and identically zero in some other part.

Lemma 2.5. *The function $f : \mathbb{R} \rightarrow \mathbb{R}$ defined by*

$$f(t) \equiv \begin{cases} e^{-1/t} & t > 0 \\ 0 & t \leq 0 \end{cases}$$

is smooth.

Proof. By induction, showing that the k th derivative of f is of the form

$$f^{(k)}(t) = \frac{p_k(t)}{t^{2k}} e^{-1/t}$$

■

Lemma 2.6. *There exists a smooth function $h : \mathbb{R} \rightarrow \mathbb{R}$, called the cutoff function, such that*

$$h(t) \equiv \begin{cases} 1 & t \leq 1 \\ 0 < h(t) < 1 & 1 < t < 2 \\ 0 & t \geq 2 \end{cases}$$

Definition 2.8. If f is any real-valued or vector-valued function on a topological space M , the *support* of f , denoted by $\text{supp} f$, is the closure of the set of points where f is nonzero.

$$\text{supp} f \equiv \text{cl}(\{p \in M \mid f(p) \neq 0\})$$

If $\text{supp} f$ is contained in some set U , we say that f is *supported in* U . A function f is said to be *compactly supported* if $\text{supp} f$ is a compact set. Clearly, every function on a compact space is compactly supported.

Lemma 2.7. *There is a smooth function $H : \mathbb{R}^n \rightarrow \mathbb{R}$ such that $0 \leq H(x) \leq 1$ everywhere, $H \equiv 1$ on $\bar{B}_1(0)$, and $\text{supp} H = \bar{B}_2(0)$, where $B_i(c)$ is the open ball with radius i centered at c .*

Proof. Just set $H(x) = h(\|x\|)$, where h is the cutoff function. ■

We can visualize the function H in the preceding lemma by assigning a greyscale color to the space \mathbb{R}^n (1 representing black, 0 representing white, and everything in between is greyscale). Then, H would produce a black closed ball of radius 1 in \mathbb{R}^n , with a smoothly changing shade of grey outside the ball of radius 1 but in the ball of radius 2, which then smoothly transitions to white for the rest of $\mathbb{R}^n \setminus B_2(0)$. The function H constructed in this lemma is an example of a *smooth bump function*, a smooth real-valued function that is equal to 1 on a specified closed set (in this case, $\bar{B}_1(0)$) and is supported in a specified open set (in this case, any open set containing $\bar{B}_2(0)$). Later, we will generalize this notion to manifolds.

2.5.1 Paracompactness

In order to rigorously define the existence of partitions of unity, we must introduce some technical definitions. The biggest takeaway from this section is that a smooth manifold M that is paracompact admits a smooth partition of unity.

Definition 2.9. Let X be a topological space. A collection \mathcal{U} of subsets of X is said to be *locally finite* if each point of X has a neighborhood that intersects at most finite many of the sets in \mathcal{U} .

Clearly, an open cover of X where each open set intersects finitely many others is locally finite.

Definition 2.10. Given an open cover \mathcal{U} of X , another open cover \mathcal{V} is called a *refinement* of \mathcal{U} if for each $V \in \mathcal{V}$ there exists some $U \in \mathcal{U}$ such that $V \subset U$ (in a way, \mathcal{V} is finer than \mathcal{U}).

Definition 2.11. A topological space X is *paracompact* if every open cover of X admits a locally finite refinement.

Proposition 2.8. *Let M be a smooth manifold. Every open cover of M has a regular refinement. In particular, M is paracompact.*

Definition 2.12. Let M be a topological space, and let $\mathcal{X} = \{X_\alpha\}_{\alpha \in A}$ be an arbitrary open cover of M . A *partition of unity subordinate to \mathcal{X}* is a collection of continuous functions $\{\psi_\alpha : M \rightarrow \mathbb{R}\}$ with the following properties.

1. $0 \leq \psi_\alpha(x) \leq 1$ for all $\alpha \in A$ and all $x \in M$.
2. $\text{supp} \psi_\alpha \subset X_\alpha$.
3. The set of supports $\{\text{supp} \psi_\alpha\}$ is locally finite.
4. $\sum_{\alpha \in A} \psi_\alpha(x) = 1$ for all $x \in M$.

A *smooth* partition of unity is one for which each of the functions ψ_α is smooth.

Theorem 2.9 (Existence of Partitions of Unity). *If M is a smooth manifold and $\mathcal{X} = \{X_\alpha\}_{\alpha \in A}$ is any open cover of M , there exists a smooth partition of unity subordinate to \mathcal{X} .*

Proposition 2.10 (Existence of Bump Functions). *Let M be a smooth manifold. For any closed set $A \subset M$ and any open set U containing A , there exists a smooth bump function for A supported in U .*

3 Tangent Vectors

In order to utilize linear approximations on smooth manifolds, we introduce the notion of a tangent space on a manifold.

3.1 Geometric Tangent Vectors

Let us assign a vector space, called a *tangent space* to each point in \mathbb{R}^n . That is, the *geometric tangent space* to \mathbb{R}^n at the point $a \in \mathbb{R}^n$ is defined

$$\mathbb{R}_a^n \equiv \{(a, v) \mid v \in \mathbb{R}^n\}$$

under the natural operations

$$\begin{aligned} v_a + w_a &\equiv (v + w)_a \\ c(v_a) &\equiv (cv)_a \end{aligned}$$

A *geometric tangent vector* in \mathbb{R}^n is an element of this space, denoted v_a , and $\mathbb{R}_a^n \simeq \mathbb{R}^n$. Note that while these tangent spaces are isomorphic, we distinguish them with the subscripts a representing the point. When looking at other sets in Euclidean space, such as $S^{n-1} \subset \mathbb{R}^n$, we can define the tangent space as the space of vectors that are orthogonal to the radial unit vector through a . But this definition is limited within the confines of Euclidean space and not for abstract manifolds. Therefore, we must utilize the concept of directional derivatives, which is provided by a tangent vector, to construct tangent spaces.

For example, a geometric tangent vector $v_a \in \mathbb{R}_a^n$ yields a map

$$D_v|_a : C^\infty(\mathbb{R}^n) \longrightarrow \mathbb{R}$$

which takes the directional derivative in the direction v at a .

$$D_v|_a f = D_v f(a) = \left. \frac{d}{dt} \right|_{t=0} f(a + tv)$$

This differential operation at the point a is linear and satisfies the product rule

$$D_v|_a (fg) = f(a)D_v|_a g + g(a)D_v|_a f$$

If $v_a = v^i e_i|_a$ in terms of the standard basis (where $e_i|_a$ is the basis of \mathbb{R}_a^n), then by the chain rule $D_v|_a f$ can be written more concretely as

$$D_v|_a f = v^i \frac{\partial f}{\partial x^i}(a)$$

Definition 3.1. If $a \in \mathbb{R}_a^n$, a linear map $X : C^\infty(\mathbb{R}^n) \longrightarrow \mathbb{R}$ is called a *derivation at a* if it satisfies the following product rule

$$X(fg) = f(a)Xg + g(a)Xf$$

Furthermore, let $T_a(\mathbb{R}^n)$ denote the set of all derivations of $C^\infty(\mathbb{R}^n)$ at a . Let us endow this set with the operations of addition and scalar multiplication

$$\begin{aligned} (X + Y)f &\equiv Xf + Yf \\ (cX)f &\equiv c(Xf) \end{aligned}$$

It can be checked that if X, Y are derivations (i.e. satisfies linearity and the product rule), then $X + Y$ and cX are also derivations, which makes $T_a(\mathbb{R}^n)$ a vector space.

Proposition 3.1. For any $a \in \mathbb{R}^n$, the map $v_a \mapsto D_v|_a$ is an isomorphism from \mathbb{R}_a^n to $T_a(\mathbb{R}^n)$.

Corollary 3.1.1. For any $a \in \mathbb{R}^n$, the n derivations

$$\left. \frac{\partial}{\partial x^1} \right|_a, \dots, \left. \frac{\partial}{\partial x^n} \right|_a, \left. \frac{\partial}{\partial x^i} \right|_a f = \frac{\partial f}{\partial x^i}(a)$$

form a basis for $T_a(\mathbb{R}^n)$, which therefore has dimension n . These basis vectors are more commonly known as the *partial derivatives of the C^∞ function f at the point a* .

Proof. Use the fact that

$$\left. \frac{\partial}{\partial x^i} \right|_a = D_{e_i}|_a$$

■

So far, given the Euclidean space \mathbb{R}^n , we have constructed this: for every $a \in \mathbb{R}^n$, the tangent space $T_a(\mathbb{R}^n)$ consists of vectors that are also derivations of $C^\infty(\mathbb{R}^n)$. In other words, we can view the tangent space at a as the vector space of linear differential operators that each take in a C^∞ function $f : \mathbb{R}^n \longrightarrow \mathbb{R}$ and outputs the directional derivative of f in direction v , evaluated at a , which is a real number. It turns out that in each tangent space at, say $a \in \mathbb{R}^n$, there exists a basis of directional derivatives (more commonly known as the partial derivatives) in direction e_i , the basis vectors, and of course, evaluated at point a .

3.1.1 Tangent Vectors on Manifolds

Before we define tangent spaces on a smooth manifold M , recall that the definition of a smooth (or C^∞) function $f : M \rightarrow \mathbb{R}$ says that there exists a smooth chart (U, φ) for M whose domain contains p and the composite function

$$f \circ \varphi^{-1} : \mathbb{R}^n \rightarrow \mathbb{R}$$

is smooth on the open subset $\varphi(U) \subset \mathbb{R}^n$. In other words, we are defining the smoothness of f really through the smoothness of $f \circ \varphi^{-1}$.

Definition 3.2. Let M be a smooth manifold and let p be a point of M . A linear map $X : C^\infty(M) \rightarrow \mathbb{R}$ is called a *derivation at p* if it satisfies

$$X(fg) = f(p)Xg + g(p)Xf$$

for all $f, g \in C^\infty(M)$. The set of all derivatives at p is a vector space called the *tangent space to M at p* , and is denoted by T_pM . An element of T_pM is called a *tangent vector at p* .

Therefore, we can think of the tangent space of point p in a smooth manifold M as the space of all directional derivatives of a smooth function

$$f \circ \varphi^{-1} : \mathbb{R}^n \rightarrow \mathbb{R}$$

evaluated at the point $\varphi^{-1}(p)$. It is good to visualize tangent vectors to an abstract smooth manifold M as arrows that are tangent to M and whose base points are attached to M at the given point. Theorems about tangent vectors must always be proved using the abstract definition in terms of derivations, but the intuition should be guided by the geometric picture.

3.2 Pushforwards

We now observe the way that tangent vectors behave under smooth maps. In the case of a smooth map between Euclidean spaces, the total derivative of the map at a point (represented by its Jacobian matrix) is a linear map that represents the "best linear approximation" to the map near the given point. In the manifold case, there is a similar linear map, but it acts between tangent spaces.

Definition 3.3. Given smooth manifolds M and N with a smooth map $F : M \rightarrow N$, we can define for each $p \in M$ a map

$$F_* : T_pM \rightarrow T_{F(p)}N, \quad (F_*X)(f) \equiv X(f \circ F)$$

to be the *pushforward associated with F* . Note that X is a tangent vector of M , while F_*X is a tangent vector of N . Furthermore, if $f \in C^\infty(N)$, then $f \circ F \in C^\infty(M)$, which is consistent with the operations. The operation F_*X is clearly linear and it satisfies the product rule because

$$\begin{aligned} (F_*X)(fg) &= X((fg) \circ F) \\ &= X((f \circ F)(g \circ F)) \\ &= f \circ F(p)X(g \circ F) + g \circ F(p)X(f \circ F) \\ &= f(F(p))(F_*X)(g) + g(F(p))(F_*X)(f) \end{aligned}$$

Lemma 3.2 (Properties of Pushforwards). *Let $F : M \rightarrow N$ and $G : N \rightarrow P$ be smooth maps, and let $p \in M$. Then,*

1. $F_* : T_pM \rightarrow T_{F(p)}N$ is linear.
2. $(G \circ F)_* = G_* \circ F_* : T_pM \rightarrow T_{G \circ F(p)}P$.
3. $(\text{Id}_M)_* = \text{Id}_{T_pM} : T_pM \rightarrow T_pM$.
4. If f is a diffeomorphism, then $F_* : T_pM \rightarrow T_{F(p)}N$ is an isomorphism.

Note that while the tangent space is defined in terms of smooth functions on the whole manifold, coordinate charts are in general defined on open subsets. The key point is that the tangent space is really a purely local construction.

Proposition 3.3. *Suppose M is a smooth manifold with $p \in M$ and $X \in T_p M$. If f and g are both smooth functions on M that agree on some neighborhood of p , then $Xf = Xg$.*

Clearly, this implies that tangent spaces of open submanifolds can be naturally identified with those of the whole manifold.

Proposition 3.4. *Let M be a smooth manifold and let $U \subset M$ be an open submanifold with $i : U \rightarrow M$ the canonical inclusion map. Then, for any $p \in U$,*

$$i_* : T_p U \rightarrow T_p M$$

is an isomorphism.

This identification of X to $i_* X$ just says that they are the *same derivation*, with the exception that X acts on functions on the bigger manifold M instead of functions on U . This is a harmless claim, since functions can be defined locally. This means that any tangent vector $X \in T_p M$ can be unambiguously applied to functions defined only in a neighborhood of p and not necessarily on all of M .

Note that every finite-dimensional vector space has a natural smooth manifold structure that is independent of any choice of basis or norm. The follow proposition shows that the tangent space to a vector space can be naturally identified with the vector space itself.

Proposition 3.5 (Tangent Space to a Vector Space). *For each finite dimensional vector space V and each point $a \in V$, there is a natural isomorphism $V \rightarrow T_a V$ such that for any linear map $L : V \rightarrow W$ the following diagram commutes.*

$$\begin{array}{ccc} V & \xrightarrow{\cong} & T_a V \\ \downarrow L & & \downarrow L_* \\ W & \xrightarrow{\cong} & T_{L_a} W \end{array}$$

3.3 Computations in Coordinates

The work that we have done is quite abstract, so we will do some down-to-earth computations in local coordinates. Let (U, φ) be a smooth coordinate chart on M . Note that φ is a diffeomorphism from U to an open subset $\tilde{U} \subset \mathbb{R}^n$. Thus, treating $\varphi(U) = \tilde{U}$ as a submanifold of the manifold \mathbb{R}^n , the pushforward of the function $\varphi : U \rightarrow \mathbb{R}^n$

$$\varphi_* : T_p M \rightarrow T_{\varphi(p)} \mathbb{R}^n$$

is an isomorphism. We know that $T_{\varphi(p)} \mathbb{R}^n$ has a basis consisting of the derivations $\partial/\partial x^i|_{\varphi(p)}$, $i = 1, 2, \dots, n$. Therefore, the pushforwards of these vectors under $(\varphi^{-1})_*$ form a basis for $T_p M$. The images of these vectors will be denoted

$$\frac{\partial}{\partial x^i} \Big|_p = (\varphi^{-1})_* \frac{\partial}{\partial x^i} \Big|_{\varphi(p)}$$

It is clear that $\partial/\partial x^i|_p$ acts on smooth functions $f : U \rightarrow \mathbb{R}$ by

$$\frac{\partial}{\partial x^i} \Big|_p \equiv \frac{\partial}{\partial x^i} \Big|_{\varphi(p)} (f \circ \varphi^{-1}) = \frac{\partial \hat{f}}{\partial x^i} (\hat{p})$$

where $\hat{f} = f \circ \varphi^{-1}$ is the coordinate representation of f , and $\hat{p} = (p^1, \dots, p^n) = \varphi(p)$ is the coordinate representation of p . This can be summarized in the following lemma.

Lemma 3.6. *Let M be a smooth manifold. For any $p \in M$, $T_p M$ is an n -dimensional vector space. If $(U, (x^i))$ is any smooth chart containing p , the coordinate vectors*

$$\left\{ \frac{\partial}{\partial x^1} \Big|_p, \dots, \frac{\partial}{\partial x^n} \Big|_p \right\}$$

form a basis for $T_p M$. Thus any tangent vector can be written uniquely as a linear combination.

$$X = X^i \frac{\partial}{\partial x^i} \Big|_p$$

using the summation convention.

The numbers X^i are the *components* of X . If X is known, its components can be computed easily from its action on the coordinate functions. That is, for each j , we can think of x^j (which outputs the j th component of a vector) as a smooth real valued function on U , to get

$$X(x^j) = \left(X^i \frac{\partial}{\partial x^i} \Big|_p \right) (x^j) = X^i \frac{\partial x^j}{\partial x^i} (p) = X^j$$

Abridged, the components of X are given by $X^j = X(x^j)$.

Now, we observe how pushforwards look in coordinates, starting off with maps between Euclidean spaces. Let $F : U \subset \mathbb{R}^n \rightarrow V \subset \mathbb{R}^m$ be a smooth map (U, V open sets in their respective spaces). For any $p \in U$, the pushforward is a linear map $F_* : T_p \mathbb{R}^n \rightarrow T_{F(p)} \mathbb{R}^m$, which has a certain matrix representation under the standard basis coordinates. Taking a typical basis vector $\partial/\partial x^i \Big|_p$ in $T_p \mathbb{R}^n$, we find its image in $T_{F(p)} \mathbb{R}^m$ under F_* . Using the chain rule,

$$\begin{aligned} \left(F_* \frac{\partial}{\partial x^i} \Big|_p \right) f &= \frac{\partial}{\partial x^i} \Big|_p (f \circ F) = \frac{\partial f}{\partial y^j} (F(p)) \frac{\partial F^j}{\partial x^i} (p) \\ &= \left(\frac{\partial F^j}{\partial x^i} (p) \frac{\partial}{\partial y^j} \Big|_{F(p)} \right) f \end{aligned}$$

Thus,

$$F_* \frac{\partial}{\partial x^i} \Big|_p = \frac{\partial F^j}{\partial x^i} (p) \frac{\partial}{\partial y^j} \Big|_{F(p)}$$

This means that the matrix of the pushforward F_* in terms of standard coordinate bases is precisely the Jacobian matrix, or total derivative, of F .

$$\begin{pmatrix} \frac{\partial F^1}{\partial x^1} (p) & \cdots & \frac{\partial F^1}{\partial x^n} (p) \\ \vdots & \ddots & \vdots \\ \frac{\partial F^m}{\partial x^1} (p) & \cdots & \frac{\partial F^m}{\partial x^n} (p) \end{pmatrix}$$

Therefore, $F_* : T_p \mathbb{R}^n \rightarrow T_{F(p)} \mathbb{R}^m$ corresponds to the total derivative $DF(p) : \mathbb{R}^n \rightarrow \mathbb{R}^m$. Now considering the more general case of smooth maps between smooth manifolds: $F : M \rightarrow N$, the derivation is not very different. Choosing smooth coordinate charts (U, φ) for M near p and (V, ψ) for N near $F(p)$, we obtain the coordinate representation

$$\hat{F} \equiv \psi \circ F \circ \varphi^{-1} : \varphi(U \cap F^{-1}(V)) \rightarrow \psi(V)$$

Using the fact that $F \circ \varphi^{-1} = \psi^{-1} \circ \hat{F}$, we compute $F_* : T_p M \rightarrow T_{F(p)} N$

$$\begin{aligned} F_* \frac{\partial}{\partial x^i} \Big|_p &= F_* \left((\varphi^{-1})_* \frac{\partial}{\partial x^i} \Big|_{\varphi(p)} \right) \\ &= (\psi^{-1})_* \left(\hat{F}_* \frac{\partial}{\partial x^i} \Big|_{\varphi(p)} \right) \\ &= (\psi^{-1})_* \left(\frac{\partial \hat{F}^j}{\partial x^i} (\hat{p}) \frac{\partial}{\partial y^j} \Big|_{\hat{F}(\varphi(p))} \right) \\ &= \frac{\partial \hat{F}^j}{\partial x^i} (\hat{p}) \frac{\partial}{\partial y^j} \Big|_{F(p)} \end{aligned}$$

Thus, $F_* : T_p M \rightarrow T_{F(p)} N$ is represented in terms of the coordinate bases by the Jacobian matrix of the coordinate representation of F . One can see that the concept of the pushforward allows us to abstractify the Jacobian matrix into an abstract linear transformation from $T_p M$ to $T_{F(p)} N$. Because of this, the

pushforward of a smooth map $F : M \rightarrow N$ is sometimes called its differential or its total derivative. We will denote F_* for the pushforward of a smooth map between smooth manifolds and $DF(p)$ for the total derivative of a map between finite dimensional vector spaces.

3.3.1 Change of Coordinates

Let (U, φ) and (V, ψ) be two smooth charts on M with $p \in U \cap V$. Let us denote the coordinate function of φ by (x^i) and those of ψ by (\tilde{x}^i) . Then, any tangent vector at p can be represented to either basis $(\partial/\partial x^i|_p)$ or $(\partial/\partial \tilde{x}^i|_p)$.

It isn't too difficult to find out how these two representations are related. With the transition map

$$\psi \circ \varphi^{-1} : \varphi(U \cap V) \rightarrow \psi(U \cap V), \quad \psi \circ \varphi^{-1}(x) = (\tilde{x}^1(x), \dots, \tilde{x}^n(x))$$

the pushforward of $\psi \circ \varphi^{-1} \in \text{End}(T_p \mathbb{R}^n)$ can be written

$$(\psi \circ \varphi^{-1})_* \frac{\partial}{\partial x^i} \Big|_{\varphi(p)} = \frac{\partial \tilde{x}^j}{\partial x^i}(\varphi(p)) \frac{\partial}{\partial \tilde{x}^j} \Big|_{\psi(p)}$$

Letting $\hat{p} = \varphi(p)$, we compute

$$\begin{aligned} \frac{\partial}{\partial x^i} \Big|_p &= (\varphi^{-1})_* \frac{\partial}{\partial x^i} \Big|_{\varphi(p)} = (\varphi^{-1})_* (\psi \circ \varphi^{-1})_* \frac{\partial}{\partial x^i} \Big|_{\varphi(p)} \\ &= (\psi^{-1})_* \frac{\partial \tilde{x}^j}{\partial x^i}(\varphi(p)) \frac{\partial}{\partial \tilde{x}^j} \Big|_{\psi(p)} = \frac{\partial \tilde{x}^j}{\partial x^i}(\varphi(p)) (\psi^{-1})_* \frac{\partial}{\partial \tilde{x}^j} \Big|_{\psi(p)} \\ &= \frac{\partial \tilde{x}^j}{\partial x^i}(\hat{p}) \frac{\partial}{\partial \tilde{x}^j} \Big|_p \end{aligned}$$

This formula conveniently, looks exactly like the multivariate chain rule of partial derivatives in \mathbb{R}^n . Applying this to the components of a vector,

$$\begin{aligned} X &= X^i \frac{\partial}{\partial x^i} \Big|_p = X^i \left(\frac{\partial \tilde{x}^j}{\partial x^i}(\hat{p}) \frac{\partial}{\partial \tilde{x}^j} \Big|_p \right) \\ &= \left(X^i \frac{\partial \tilde{x}^j}{\partial x^i}(\hat{p}) \right) \frac{\partial}{\partial \tilde{x}^j} \Big|_p = \tilde{X}^j \frac{\partial}{\partial \tilde{x}^j} \Big|_p \end{aligned}$$

We can find that the components transform by the rule

$$\tilde{X}^j = \frac{\partial \tilde{x}^j}{\partial x^i}(\hat{p}) X^i$$

3.4 Tangent Vectors to Curves

Definition 3.4. We define a *curve* in manifold M to be a continuous map

$$\gamma : J \rightarrow M$$

where $J \in \mathbb{R}$ is an interval.

Our construction of tangent vectors and pushforwards leads to a very natural definition of tangent vectors of curves in manifolds.

Definition 3.5. If γ is a smooth curve in smooth manifold M , the *tangent vector* to γ at $t_0 \in J$ is the vector

$$\gamma'(t_0) \equiv \gamma_* \left(\frac{d}{dt} \Big|_{t_0} \right) \in T_{\gamma(t_0)} M$$

where $d/dt|_{t_0}$ is the standard coordinate basis vector for the 1 dimensional $T_{t_0} \mathbb{R}$.

The tangent vector acts on $C^\infty(M)$ functions f (that is, $f : M \rightarrow \mathbb{R}$) by

$$\gamma'(t_0)f = \left(\gamma_* \frac{d}{dt} \Big|_{t_0} \right) f = \frac{d}{dt} \Big|_{t_0} (f \circ \gamma) = \frac{d(f \circ \gamma)}{dt}(t_0)$$

where $f \circ \gamma : \mathbb{R} \rightarrow \mathbb{R}$. In other words, $\gamma'(t_0)$ is the derivation at $\gamma(t_0)$ obtained by taking the derivative of a function along γ . Note also that the path function γ is also a smooth map between manifolds.

Let (U, φ) be a smooth chart with coordinate functions (x^i) . If $\gamma(t_0) \in U$, we can write the coordinate representation of γ as $\gamma(t) = (\gamma^1(t), \dots, \gamma^n(t))$ for t near t_0 . (Note that explicitly speaking, we are really writing the shorthand form of $(\varphi \circ \gamma)(t) = ((\varphi \circ \gamma)^1(t), \dots, (\varphi \circ \gamma)^n(t))$). Then, the formula for the pushforward in coordinates becomes

$$\gamma'(t_0) = (\gamma^i)'(t_0) \frac{\partial}{\partial x^i} \Big|_{\gamma(t_0)}$$

This means that $\gamma'(t_0)$ is given by essentially the same formula it would be in Euclidean space: It is the tangent vector whose components in a coordinate basis are the derivatives of the component functions of γ .

Lemma 3.7. *Let M be a smooth manifold and $p \in M$. Then, every $X \in T_p M$ is the tangent vector to some smooth curve in M .*

This lemma tells us that we can think of the tangent space at p as the set of tangent vectors to smooth curves in M passing through p . Similar results hold under compositions of smooth maps.

Proposition 3.8. *Let $F : M \rightarrow N$ be a smooth map, and let $\gamma : J \rightarrow M$ be a smooth curve. For any $t_0 \in J$, the tangent vector at $t = t_0$ to the composite curve $F \circ \gamma : J \rightarrow N$ is given by*

$$(F \circ \gamma)'(t_0) = F_*(\gamma'(t_0))$$

We can think of this composition as the double pushforward of the basis vector $d/dt|_{t_0}$ to $T_{\gamma(t_0)}M$ first and $T_{(F \circ \gamma)(t_0)}N$ second).

4 Vector Fields

4.1 The Tangent Bundle

Definition 4.1. For any smooth manifold M , we define the *tangent bundle* of M , denoted TM to be the disjoint union of the tangent spaces at all points of M .

$$TM \equiv \bigsqcup_{p \in M} T_p M$$

Elements of TM are written as an ordered pair

$$(p, X) \text{ or } X_p, \quad p \in M, X \in T_p M$$

The tangent bundle also comes with an natural *projection map*

$$\pi : TM \rightarrow M, \quad \pi(p, X) = p$$

which sends each vector in $T_p M$ to the point p at which it is tangent.

The following lemma reveals an important property of tangent bundles.

Lemma 4.1. *For any smooth n -manifold M , the tangent bundle TM has a natural topology and smooth structure that make it into a $2n$ -dimensional smooth manifold. With this structure,*

$$\pi : TM \rightarrow M$$

is a smooth map.

4.2 Vector Fields on Manifolds

Definition 4.2. Given a smooth manifold M , a *vector field on M* is a continuous map

$$Y : M \longrightarrow TM, \quad p \mapsto Y_p$$

with the property that

$$\pi \circ Y = \text{Id}_M$$

or equivalently, $Y_p \in T_pM$ for each $p \in M$. We also say that the vector field is a *section* (as in *cross section*) of the map $\pi : TM \longrightarrow M$ (i.e. a continuous right inverse of π).

While each vector in the vector field Y "lives" in distinct tangent spaces, we can visualize a vector field on manifold M as an arrow attached to each point of M , chosen to be tangent to M and to vary continuously from point to point.

Definition 4.3. A *smooth vector field* is a smooth map from M to TM . A *rough vector field* is a (not necessarily continuous) map $Y : M \longrightarrow TM$ satisfying the condition that $\pi \circ Y = \text{Id}_M$.

If $Y : M \longrightarrow TM$ is a rough vector field and $(U, (x^i))$ is any smooth coordinate chart for M , we can write the value of Y at any point $p \in U$ in terms of the coordinate basis vectors.

$$Y_p = Y^i(p) \left. \frac{\partial}{\partial x^i} \right|_p$$

This defines n functions $Y^i : U \longrightarrow \mathbb{R}$, called the *component functions* of Y in the given chart.

Lemma 4.2 (Smoothness Criterion for Vector Fields). *Let M be a smooth manifold, and let $Y : M \longrightarrow TM$ be a rough vector field. If $(U, (x^i))$ is any smooth coordinate chart on M , then Y is smooth on U if and only if its component functions with respect to this chart are smooth.*

With this lemma, it is now sufficient to check the smoothness of the component functions in order to check the smoothness of the entire vector field. The next lemma shows that every tangent vector at a point can be extended to a smooth global vector field.

Lemma 4.3. *Let M be a smooth manifold. If $p \in M$ and $X \in T_pM$, then there exists a smooth vector field \tilde{X} on M such that $\tilde{X}_p = X$.*

Definition 4.4. Just as for functions, the *support* of a vector field Y is defined to be the closure of the set $\{p \in M \mid Y_p \neq 0\}$. A vector field is said to be *compactly supported* if its support is a compact set.

Furthermore, if U is any open set of M , the fact that T_pU is naturally identified with T_pM for each $p \in U$ allows us to identify the subset $\pi^{-1}(U) \subset TM$. Therefore, a vector field on U can be thought of either as a map from U to TU or a map from U to TM . If Y is a vector field on M , its restriction $Y|_U$ is a vector field on U , which is smooth if Y is.

Definition 4.5. Let us denote the set of all smooth vector fields on M as $\mathcal{T}(M)$. It is a vector space under pointwise addition and scalar multiplication. That is, given $Y, Z \in \mathcal{T}(M)$

$$(aY + bZ)_p = aY_p + bZ_p$$

The zero element of this vector space is also the zero vector field. In addition, smooth vector fields can be multiplied by smooth real-valued functions. That is, if $f \in C^\infty(M)$ and $Y \in \mathcal{T}(M)$, then we can define $fY : M \longrightarrow TM$ as

$$(fY)_p = f(p)Y_p$$

The coordinate representation of a vector field Y can also be written as an equation between vector fields rather than at a single point.

$$Y_p = Y^i(p) \left. \frac{\partial}{\partial x^i} \right|_p \implies Y = Y^i \frac{\partial}{\partial x^i}$$

A property of vector fields is that they induce operators on the space of smooth real-valued functions $C^\infty(M)$. That is, if $Y \in \mathcal{T}(M)$ and $f \in C^\infty(U)$, where U is an open set in M , then we obtain a new function $Yf : U \rightarrow \mathbb{R}$, defined

$$Yf \equiv Y^i \frac{\partial}{\partial x^i}(f) \implies Yf(p) \equiv Y_p f \equiv Y^i(p) \frac{\partial}{\partial x^i} f \Big|_p$$

Note that there is a concrete difference between Yf and fY . Yf is a real valued function shown above, while fY is a smooth vector field gotten by multiplying the vector produced by Y at point p (Y_p) with the value of f at p ($f(p)$). Because the action of a tangent vector on a function is determined by the values of the function in an arbitrarily small neighborhood, it follows that Yf is locally determined. This leads to another sufficient condition for smoothness of vector fields.

Lemma 4.4. *Let M be a smooth manifold, and let $Y : M \rightarrow TM$ be a rough vector field. Then, Y is smooth if and only if for every open set $U \subset M$ and every $f \in C^\infty(U)$, the function $Yf : U \rightarrow \mathbb{R}$ is smooth.*

Notice that due to this lemma, a smooth vector field $Y \in \mathcal{T}(M)$ defines a map from $C^\infty(M)$ to itself by the mapping $f \mapsto Yf$. This is clearly linear over \mathbb{R} and satisfies the product rule

$$Y(fg) = fYg + gYf$$

since it satisfies for arbitrary point $p \in M$.

Definition 4.6. A linear endomorphism Y of $C^\infty(M)$ satisfying

$$Y(fg) = fYg + gYf, \quad f, g \in C^\infty(M)$$

is called a *derivation* of $C^\infty(M)$.

The next proposition shows that derivations of $C^\infty(M)$ can be identified with smooth vector fields.

Proposition 4.5. *Let M be a smooth manifold. A map $\mathcal{Y} : C^\infty(M) \rightarrow C^\infty(M)$ is a derivation if and only if it is of the form $\mathcal{Y}f = Yf$ for some smooth vector field $Y \in \mathcal{T}(M)$.*

4.2.1 Pushforwards of Vector Fields

If $F : M \rightarrow N$ is a smooth map and Y is a vector field on M , then for each point $p \in M$, we obtain a vector $F_*Y_p \in T_{F(p)}N$ by pushing forward Y_p . However, this does not in general define a vector field on N . For example, if F is not surjective, we cannot assign a vector to point $q \in N \setminus F(M)$. If F is not injective, then for some point of N there may be several different vectors obtained as pushforwards of Y from different points of M . Therefore, vector fields do not always push forward.

Definition 4.7. If $F : M \rightarrow N$ is smooth and Y is a vector field on M , suppose there happens to be a vector Z on N with the property that for each $p \in M$, $F_*Y_p = Z_{F(p)}$. In this case, we say that the vector fields Y and Z are *F-related*.

The vector field Z in the definition above represents the "closest" vector field on N that we can get that is a pushforward of Y . If F is not surjective, the vector field Z can take any value at points $p \in N \setminus F(M)$ (meaning that the F -related vector field to Y is not unique). If F is not injective, for all points a_i that map onto $r \in N$, the pushforward of F at each a_i must agree on their output vector in T_rN . There is a useful criterion to see if two vector fields are F -related.

Proposition 4.6. *Suppose $F : M \rightarrow N$ is a smooth map, $Y \in \mathcal{T}(M)$, and $Z \in \mathcal{T}(N)$. Then, Y and Z are F -related if and only if for every smooth real-valued function defined on an open subset of N ,*

$$Y(f \circ F) = (Zf) \circ F$$

Note that for a given smooth map $F : M \rightarrow N$ and vector field $Y \in \mathcal{T}(M)$, there may not be *any* vector field on N that is F -related to Y . However, there is one special case in which there always exists a unique vector field.

Proposition 4.7. *Suppose $F : M \rightarrow N$ is a diffeomorphism. For every $Y \in \mathcal{T}(M)$, there is a unique smooth vector field on N that is F -related to Y .*

It is quite easy to see why the proposition above is true. Since F is a diffeomorphism (and thus a bijection), there will be no points on N where the F -related vector field Z is undefined (due to surjectivity) and there will be no contradictions in the values of Z (due to injectivity). Finally, F being a diffeomorphism will ensure smoothness of Z .

Definition 4.8. In the case where $F : M \rightarrow N$ is a diffeomorphism, the unique vector field F_*Y that is F -related to Y is called the *pushforward of Y by F* . Note that F_*Y is defined only if F is a diffeomorphism.

4.2.2 Vector Fields on Manifolds with Boundary

4.3 Lie Brackets

We can fix this problem using the Lie bracket operator.

Definition 4.9. The operator

$$[\cdot, \cdot] : \mathcal{T}(M) \times \mathcal{T}(M) \rightarrow \mathcal{T}(M)$$

is called the *Lie bracket*. Given two vector fields V, W on smooth manifold M , we define the $[V, W]$ as

$$[V, W] : C^\infty(M) \rightarrow C^\infty(M), \quad [V, W]f \equiv VWf - WVf$$

Lemma 4.8. *The Lie bracket of any pair of smooth vector fields is a smooth vector field.*

The value of the vector field $[V, W]$ at a point $p \in M$ is the derivation at p given by the formula

$$[V, W]_p f \equiv V_p(Wf) - W_p(Vf)$$

but the formula is of limited usefulness for practical computations. The following lemma simplifies it greatly.

Lemma 4.9. *Let V, W be smooth vector fields on a smooth manifold M , and let*

$$V = V^i \frac{\partial}{\partial x^i} \text{ and } W = W^j \frac{\partial}{\partial x^j}$$

be the coordinate expressions for V and W in terms of some smooth local coordinates (x^i) for M . Then, $[V, W]$ has the following coordinate expression:

$$[V, W] = \left(V^i \frac{\partial W^j}{\partial x^i} - W^i \frac{\partial V^j}{\partial x^i} \right) \frac{\partial}{\partial x^j}$$

or more concisely,

$$[V, W] = (VW^j - WV^j) \frac{\partial}{\partial x^j}$$

Proof. Since $[V, W]$ is a smooth vector field, its values are determined locally. Thus, it suffices to compute in a single smooth chart.

$$\begin{aligned} [V, W]f &= V^i \frac{\partial}{\partial x^i} \left(W^j \frac{\partial f}{\partial x^j} \right) - W^j \frac{\partial}{\partial x^j} \left(V^i \frac{\partial f}{\partial x^i} \right) \\ &= V^i \frac{\partial W^j}{\partial x^i} \frac{\partial f}{\partial x^j} + V^i W^j \frac{\partial^2 f}{\partial x^i \partial x^j} - W^j \frac{\partial V^i}{\partial x^j} \frac{\partial f}{\partial x^i} - W^j V^i \frac{\partial^2 f}{\partial x^j \partial x^i} \\ &= V^i \frac{\partial W^j}{\partial x^i} \frac{\partial f}{\partial x^j} - W^j \frac{\partial V^i}{\partial x^j} \frac{\partial f}{\partial x^i} \end{aligned}$$

where we've used the product rule in the first to second step and the fact that mixed partial derivatives of a smooth function are equal in any order. ■

Example 4.1. *Let us define smooth vector fields $V, W \in \mathcal{T}(\mathbb{R}^3)$ by*

$$\begin{aligned} V &= x \frac{\partial}{\partial x} + \frac{\partial}{\partial y} + x(y+1) \frac{\partial}{\partial z} \\ W &= \frac{\partial}{\partial x} + y \frac{\partial}{\partial z} \end{aligned}$$

After some tedious calculations, we compute a total of nine separate terms to get

$$\begin{aligned} [V, W] &= -\frac{\partial}{\partial x} - (y+1)\frac{\partial}{\partial z} + \frac{\partial}{\partial z} \\ &= -\frac{\partial}{\partial x} - y\frac{\partial}{\partial z} \end{aligned}$$

Lemma 4.10 (Properties of the Lie Bracket). *The Lie bracket satisfies the following identities for all $V, W, X \in \mathcal{T}(M)$.*

1. *Bilinearity.* For all $a, b \in \mathbb{R}$,

$$\begin{aligned} [aV + bW, X] &= a[V, X] + b[W, X] \\ [X, aV + bW] &= a[X, V] + b[X, W] \end{aligned}$$

2. *Antisymmetry.*

$$[V, W] = -[W, V]$$

3. *Jacobi Identity.*

$$[V, [W, X]] + [W, [X, V]] + [X, [V, W]] = 0$$

4. For $f, g \in C^\infty(M)$,

$$[fV, gW] = fg[V, W] + (fVg)W - (gWf)V$$

The next proposition states the naturality of the Lie bracket. That is, F -relatedness is preserved with the Lie bracket.

Proposition 4.11. *Let $F : M \rightarrow N$ be a smooth map, and let $V_1, V_2 \in \mathcal{T}(M)$ and $W_1, W_2 \in \mathcal{T}(N)$ be vector fields such that V_i is F -related to W_i for $i = 1, 2$. Then, $[V_1, V_2]$ is F -related to $[W_1, W_2]$.*

Corollary 4.11.1. *Suppose $F : M \rightarrow N$ is a diffeomorphism and $V_1, V_2 \in \mathcal{T}(M)$. Then*

$$F_*[V_1, V_2] = [F_*V_1, F_*V_2]$$

4.4 The Lie Algebra of a Lie Group

Definition 4.10. Let G be a Lie group. Any $g \in G$ defines maps $L_g, R_g : G \rightarrow G$, called *left translation* and *right translation*, respectively, by

$$L_g(h) \equiv gh, R_g(h) \equiv hg$$

Because L_g can be written as the composition of smooth maps

$$G \xrightarrow{i_g} G \times G \xrightarrow{m} G$$

where $i_g(h) \equiv (g, h)$ and m is multiplication, it follows that L_g is smooth. It is actually a diffeomorphism of G since $L_{g^{-1}}$ is a smooth inverse for it. The same goes for R_g . Notice that given any two points $g_1, g_2 \in G$, there is a unique left translation of G taking g_1 to g_2 . This is the translation $g_2g_1^{-1}$. Many important Lie groups follow from the fact that we can systematically map any point to any other by such a global diffeomorphism.

Definition 4.11. A vector field X on G is said to be *left-invariant* if it is invariant under all left translation, in the sense that it is L_g -related to itself for every $g \in G$. More explicitly, this means that

$$(L_g)_*X_{g'} = X_{gg'} \text{ for all } g, g' \in G$$

Furthermore, because

$$(L_g)_*(aX + bY) = a(L_g)_*X + b(L_g)_*Y$$

the set of all smooth left-invariant vector fields on G is a linear subspace of $\mathcal{T}(M)$. Even further, it is a *Lie algebra*, which will be defined shortly.

Lemma 4.12. *Let G be a Lie group, and suppose X and Y are smooth left-invariant vector fields on G . Then $[X, Y]$ is left-invariant.*

Proof. Since $(L_g)_*X = X$ and $(L_g)_*Y = Y$ by definition of left-invariance, it follows that

$$(L_g)_*[X, Y] = [(L_g)_*X, (L_g)_*Y] = [X, Y]$$

meaning that $[X, Y]$ is L_g -related to itself. ■

Definition 4.12. A *Lie algebra* is a real vector space \mathfrak{g} endowed with a map called the *bracket* from $\mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$, usually denoted by $(X, Y) \mapsto [X, Y]$, that satisfies the following properties for all $X, Y, Z \in \mathfrak{g}$.

1. Bilinearity. For all $a, b \in \mathbb{R}$,

$$\begin{aligned} [aV + bW, X] &= a[V, X] + b[W, X] \\ [X, aV + bW] &= a[X, V] + b[X, W] \end{aligned}$$

2. Antisymmetry.

$$[V, W] = -[W, V]$$

3. Jacobi Identity.

$$[V, [W, X]] + [W, [X, V]] + [X, [V, W]] = 0$$

Notice that the Jacobi identity is a substitute for associativity, which does not hold in general for brackets in a Lie algebra.

Definition 4.13. If \mathfrak{g} is a Lie algebra, a linear subspace $\mathfrak{h} \subset \mathfrak{g}$ is called a *Lie subalgebra* of \mathfrak{g} if it is closed under brackets. In other words, \mathfrak{h} is itself a Lie algebra under the restriction of the bracket to \mathfrak{h} .

Definition 4.14. If \mathfrak{g} and \mathfrak{h} are Lie algebras, a linear map

$$A : \mathfrak{g} \rightarrow \mathfrak{h}$$

if it preserves brackets. That is, if

$$A[X, Y] = [AX, AY]$$

An invertible Lie algebra homomorphism is called a *Lie algebra isomorphism*. Two Lie algebra with an isomorphism between them are said to be *isomorphic Lie algebras*.

Example 4.2. *The space $\mathcal{T}(M)$ of all smooth vector fields on a smooth manifold M is a Lie algebra under the Lie bracket.*

Example 4.3. *Any vector space V becomes a Lie algebra if we define all brackets to be 0. Such a Lie algebra is said to be abelian, because the underlying product \cdot in the commutator $[A, B] = A \cdot B - B \cdot A$ is commutative.*

Example 4.4. *If V is a vector space, the linear space of endomorphisms of V , denoted $\mathfrak{gl}(V)$, becomes a Lie algebra with the commutator bracket*

$$[A, B]x \equiv A(Bx) - B(Ax)$$

This can be represented using matrices, making the set of all $n \times n$ matrices with the commutator defined $[A, B] \equiv AB - BA$ also a Lie algebra.

Definition 4.15. If G is a Lie group, the set of all smooth left-invariant vector fields on G is a Lie subalgebra of $\mathcal{T}(G)$ and it therefore a Lie algebra. This Lie algebra is denoted $\text{Lie}(G)$.

Note that we have already proved that the set of smooth left-invariant vector fields on G is closed under the Lie bracket.

Theorem 4.13. *Let G be a Lie group. The evaluation map*

$$\varepsilon : \text{Lie}(G) \rightarrow T_e G, \quad \varepsilon(X) = X_e$$

where e is the identity element of G , is a vector space isomorphism. Thus, $\text{Lie}(G)$ is finite dimensional, with dimension equal to $\dim G$.

Proof. Will be done. ■

Note also that the preceding proof shows that the assumption of smoothness in the definition of $\text{Lie}(G)$ is unnecessary.

Corollary 4.13.1. *Every left-invariant rough vector field on a Lie group is smooth.*

Proposition 4.14 (Lie algebra of the General Linear Group). *The composition of the natural maps*

$$\text{Lie}(\text{GL}(n, \mathbb{R})) \rightarrow T_{I_n} \text{GL}(n, \mathbb{R}) \rightarrow \mathfrak{gl}(n, \mathbb{R})$$

gives a Lie algebra isomorphism between $\text{Lie}(\text{GL}(n, \mathbb{R}))$ and the matrix algebra $\mathfrak{gl}(n, \mathbb{R})$.

4.4.1 Induced Lie Algebra Homomorphisms

It can be seen that each Lie group homomorphism induces a Lie algebra homomorphism.

Theorem 4.15. *Let G and H be Lie groups, and let \mathfrak{g} and \mathfrak{h} be their Lie algebras. Suppose $F : G \rightarrow H$ is a Lie group homomorphism. For every $X \in \mathfrak{g}$, there is a unique vector field \mathfrak{h} that is F -related to X . With this vector field, denoted F_*X , the map $F_* : \mathfrak{g} \rightarrow \mathfrak{h}$ so defined is a Lie algebra homomorphism.*

Lemma 4.16 (Properties of the Induced Homomorphism). *1. The homomorphism $(\text{Id}_G)_* : \text{Lie}(G) \rightarrow \text{Lie}(G)$ induced by the identity map of G is the identity of $\text{Lie}(G)$.*

2. If $F_1 : G \rightarrow H$ and $F_2 : H \rightarrow K$ are Lie group homomorphisms, then $(F_2 \circ F_1)_ = (F_2)_* \circ (F_1)_* : \text{Lie}(G) \rightarrow \text{Lie}(G)$.*

3. Isomorphic Lie groups have isomorphic Lie algebras.

5 Vector (Fiber) Bundles

We have already seen that the tangent bundle of a smooth manifold has a natural structure as a smooth manifold in its own right. The standard coordinates on TM make it look, locally, like $M \times \mathbb{R}^n$. This kind of structure arises frequently. That is, a collection of vector spaces (one for each point in M) glued together so that it looks *locally* like the Cartesian product of M with \mathbb{R}^n , but globally it may not be. This is called the vector bundle. Note that a tangent bundle is one type of vector bundle.

Definition 5.1. Let M be a topological space. A (real) vector bundle of rank k over M is a topological space E together with a surjective continuous map $\pi : E \rightarrow M$ satisfying:

1. For each $p \in M$, the set $E_p \equiv \pi^{-1}(p) \subset E$ (called the *fiber of E over p*) is endowed with the structure of a k -dimensional real vector space.
2. For each $p \in M$, there exists a neighborhood U of p in M and a homeomorphism $\Phi : \pi^{-1}(U) \rightarrow U \times \mathbb{R}^k$ (called a *local trivialization of E over U*), such that the following diagram commutes:

$$\begin{array}{ccc} \pi^{-1}(U) & \xrightarrow{\Phi} & U \times \mathbb{R}^k \\ \downarrow \pi & \swarrow \pi_1 & \\ U & & \end{array}$$

where π_1 is the projection onto the first factor. Furthermore, for each $q \in U$, the restriction of Φ to E_q is a linear isomorphism from E_q to $\{q\} \times \mathbb{R}^k \simeq \mathbb{R}^k$.

Note that E is *not* the same space as $M \times \mathbb{R}^k$, but rather every two tuple in $M \times \mathbb{R}^k$ can be *associated* with every element in E through the surjective map π (unless E is a trivial bundle).

Definition 5.2. If M and E are smooth manifolds, π is a smooth projection map, and the local trivializations can be chosen to be diffeomorphisms, then E is called a *smooth vector bundle*. In this case, we will call any local trivialization that is a diffeomorphism onto its image a *smooth local trivialization*. The space E is called the *total space* of the bundle, M called its *base*, and π called its *projection*. Also, if $U \subset M$ is any open set, it is easy to verify that the subset $E|_U = \pi^{-1}(U)$ is again a vector bundle with the restriction of π as its projection map, called the *restriction of E to U* .

A rank-1 vector bundle is often called a *line bundle*. Complex bundles are also defined with vector space over \mathbb{C} .

Definition 5.3. If there exists a local trivialization over all of M (called a *global trivialization of E*), then E is said to be a *trivial bundle*. In this case, E itself is homeomorphic to the product space $M \times \mathbb{R}^k$. If $E \rightarrow M$ is a smooth bundle that admits a smooth global trivialization, then we say that E is *smoothly trivial* (in this case, E is diffeomorphic to $M \times \mathbb{R}^k$, not just homeomorphic).

Example 5.1 (Trivial Bundle). Let $E = M \times \mathbb{R}^k$, and let $\pi = \pi_1 : M \times \mathbb{R}^k \rightarrow M$ be the projection onto the first factor. Then E is a fiber bundle of \mathbb{R}^k over M . Here E is not just locally a product but globally one. Every trivial bundle is of this form, with the identity map as a global trivialization. If M is a smooth manifold, then $M \times \mathbb{R}^k$ is smoothly trivial.

Example 5.2 (The Cylinder as a Trivial Bundle). Let E be a cylinder, i.e. a cylindrical surface. We remind the reader that E stretches infinitely long, and that it is "hollow." Then the 1-sphere, denoted S^1 , is the base space, and \mathbb{R} is the fiber. Geometrically, we can think of S^1 to be the base of the cylinder, while an infinite number of real lines orthogonally intersect S^1 at one point to cover E . We can then define a map $S^1 \times \mathbb{R} \rightarrow E$ to be

$$(\theta, z) \rightarrow (\cos \theta, \sin \theta, z)$$

The standard fiber F is \mathbb{R} , but we must note that there are an infinite number of copies of \mathbb{R} passing through each point in E . So given points $p, q \in E$, the specific fibers associated with p and q are different \mathbb{R} s. We can think of this fiber \mathbb{R} as the *collection of all real lines that fill the surface of cylinder E* .

Note that we could have very well just chosen \mathbb{R} to be the base space and S^1 to be the fiber. The base space would then be unbounded, but this doesn't pose as a problem.

Furthermore, in the example shown before, the map $B \times F \rightarrow E$ doesn't have to be so simple as

$$(z, \theta) \rightarrow (\cos \theta, \sin \theta, z)$$

In this mapping, all of the \mathbb{R} passing through each point is parameterized the same, since the same value of z gives the same heights everywhere. However, for the more complex parameterization

$$(\theta, z) \rightarrow (\cos \theta, \sin \theta, (2 + \sin \theta)z)$$

the parameterizations in each \mathbb{R} are different for every point $\theta \in S^1$. So, the same value of z will not give you the same height everywhere. However, note that while all of these \mathbb{R} are parameterized differently, every single \mathbb{R} are diffeomorphic since, once picking all of them to have the same parameterization, we can identify the identity mapping as a diffeomorphism between all of them.

Usually, we deal with nontrivial bundles that have a cover of open sets. Here is an example.

Example 5.3 (Möbius Bundle). Let $I = [0, 1] \subset \mathbb{R}$ be the unit interval, and let $p : I \rightarrow S^1$ be the quotient map $p(x) \equiv e^{2\pi ix}$, which identifies the two endpoints of I . Consider the "infinite strip" $I \times \mathbb{R}$, and let $\pi : I \times \mathbb{R} \rightarrow I$ be the projection on the first factor. Let \sim be the equivalence relation on $I \times \mathbb{R}$ that identifies each point $(0, y)$ in the fiber over 0 with the point $(1, -y)$ in the fiber over 1. Geometrically, we are half-twisting the right hand side of the strip and gluing it into the left-hand edge. The resulting quotient space is $E \equiv (I \times \mathbb{R}) / \sim$, with $q : I \times \mathbb{R} \rightarrow E$ the quotient map.

Because $p \circ \pi_1$ is constant on each equivalence class, it descends to a continuous map $p \circ \pi_1 = \pi : E \rightarrow S^1$. This makes E into a smooth real line bundle over S^1 .

Proposition 5.1 (The Tangent Bundle as a Vector Bundle). Let M be a smooth n -manifold and let TM be its tangent bundle. With its standard projection map, its natural vector space structure on each fiber, and the smooth manifold structure, TM is a smooth vector bundle of rank n over M .

Proof. Given any smooth chart (U, φ) for M with coordinate functions (x^i) , we define the map $\Phi : \pi^{-1}(U) \rightarrow U \times \mathbb{R}^n$ by

$$\Phi \left(v^i \frac{\partial}{\partial x^i} \Big|_p \right) \equiv (p, (v^1, v^2, \dots, v^n))$$

This is obviously linear on fibers and satisfies $\pi \circ \Phi = \pi$. The composite map

$$\pi^{-1}(U) \xrightarrow{\Phi} U \times \mathbb{R}^n \xrightarrow{\varphi \times \text{Id}_{\mathbb{R}^n}} \varphi(U) \times \mathbb{R}^n$$

Since both $\tilde{\varphi}$ and $\varphi \times \text{Id}_{\mathbb{R}^n}$ are diffeomorphisms, Φ is also a diffeomorphism, satisfying all the conditions of a local trivialization. ■

We still have to deal with overlaps between trivializing neighborhoods. The next lemma introduces transition functions between different representations of a fiber.

Lemma 5.2 (Overlaps between Trivializing Neighborhoods). *Let $\pi : E \rightarrow M$ be a smooth vector bundle, and suppose $\Phi : \pi^{-1}(U) \rightarrow U \times \mathbb{R}^k$ and $\Psi : \pi^{-1}(V) \rightarrow U \times \mathbb{R}^k$ be two smooth local trivializations of E such that $U \cap V \neq \emptyset$. Then, there exists a smooth map $\tau : U \cap V \rightarrow \text{GL}(k, \mathbb{R})$ such that the composition $\Phi \circ \Psi^{-1} : (U \cap V) \times \mathbb{R}^k \rightarrow (U \cap V) \times \mathbb{R}^k$ has the form*

$$\Phi \circ \Psi^{-1}(p, v) \equiv (p, \tau(p)v)$$

where $\tau(p)v$ denotes the usual action of the $k \times k$ matrix $\tau(p)$ on the vector $v \in \mathbb{R}^k$. Furthermore, $\Phi \circ \Psi^{-1}$ is a diffeomorphism. We can represent this consistency by the following commutative diagram.

$$\begin{array}{ccc} (U \cap V) \times \mathbb{R}^k & \xleftarrow{\Psi} \pi^{-1}(U \cap V) \xrightarrow{\Phi} & (U \cap V) \times \mathbb{R}^k \\ & \searrow \pi_1 \quad \downarrow \pi \quad \swarrow \pi_1 & \\ & & U \cap V \end{array}$$

Definition 5.4. The smooth map $\tau : U \cap V \rightarrow \text{GL}(k, \mathbb{R})$ is called the *transition function* between the local trivializations Φ and Ψ . For example, if M is a smooth manifold and Φ, Ψ are the local trivializations of TM associated with two different smooth charts, then the transition functions between them is just the Jacobian matrix of the coordinate transition map.

In order to make a vector bundle into a smooth one, we would need to construct a manifold topology and a smooth structure on the disjoint union of all the vector spaces, and then construct the local trivializations and show that they have the requisite properties. The following lemma provides a shortcut by showing that it is sufficient to construct the local trivializations, as long as they overlap with smooth transition functions.

Lemma 5.3 (Vector Bundle Construction Lemma). *Let M be a smooth manifold and suppose that we are given for each $p \in M$, a real vector space E_p of some fixed dimension k . Furthermore, let*

$$E \equiv \bigsqcup_{p \in M} E_p$$

and let $\pi : E \rightarrow M$ be the map that takes each element of E_p to point p . Suppose furthermore that we are given

1. an open cover $\{U_\alpha\}_{\alpha \in A}$ of M .
2. for each $\alpha \in A$, a bijective map $\Phi_\alpha : \pi^{-1}(U_\alpha) \rightarrow U_\alpha \times \mathbb{R}^k$ whose restriction to each E_p is a linear isomorphism from E_p to $\{p\} \times \mathbb{R}^k \simeq \mathbb{R}^k$.
3. for each $\alpha, \beta \in A$ such that $U_\alpha \cap U_\beta \neq \emptyset$, a smooth map $\tau_{\alpha\beta} : U_\alpha \cap U_\beta \rightarrow \text{GL}(k, \mathbb{R})$ such that the composite map $\Phi_\alpha \circ \Phi_\beta^{-1}$ from $((U_\alpha \cap U_\beta) \times \mathbb{R}^k)$ to itself has the form

$$\Phi_\alpha \circ \Phi_\beta^{-1}(p, v) \equiv (p, \tau_{\alpha\beta}(p)v)$$

Then, E has a unique smooth manifold structure making it into a smooth vector bundle of rank k over M , with π as projection and the maps Φ_α as smooth local trivializations.

5.1 Categories and Functors

We summarize the basic definitions of category theory, which provides a convenient and powerful language for talking about many mathematical structures.

Definition 5.5. A *category* \mathbf{C} consists of three things:

1. a class of *objects*
2. for each pair X, Y of objects a set $\text{Hom}_{\mathbf{C}}(X, Y)$ whose elements are called *morphisms*

3. for each triple X, Y, Z of objects a map called *composition*

$$\text{Hom}_{\mathbf{C}}(X, Y) \times \text{Hom}_{\mathbf{C}}(Y, Z) \longrightarrow \text{Hom}_{\mathbf{C}}(X, Z)$$

written $(f, g) \mapsto g \circ f$

The morphisms are also required to satisfy the following properties:

1. Associativity: $(f \circ g) \circ h = f \circ (g \circ h)$
2. Existence of Identities: For each object X in \mathbf{C} , there exists an *identity morphism* $\text{Id}_X \in \text{Hom}_{\mathbf{C}}(X, X)$ satisfying

$$\text{Id}_Y \circ f = f = f \circ \text{Id}_X$$

for all $f \in \text{Hom}_{\mathbf{C}}(X, Y)$

Definition 5.6. A morphism $f \in \text{Hom}_{\mathbf{C}}(X, Y)$ is called an *isomorphism* in \mathbf{C} if there exists a morphism $g \in \text{Hom}_{\mathbf{C}}(Y, X)$ such that $f \circ g = \text{Id}_Y$ and $g \circ f = \text{Id}_X$.

Example 5.4 (Categories). *When working with categories, the objects are sets with some extra structure, the morphisms are maps that preserve that structure, and the composition laws and identity morphisms are the obvious ones. Some common examples are:*

1. SET: Sets and maps
2. TOP: Topological spaces and continuous maps.
3. TM: Topological manifolds and continuous maps.
4. SM: Smooth manifolds and smooth maps.
5. VB: Smooth vector bundles and smooth bundle maps.
6. VECT $_{\mathbb{R}}$: Real vector spaces and real-linear maps.
7. VECT $_{\mathbb{C}}$: Complex vector spaces and complex-linear maps.
8. GROUP: Groups and group homomorphisms.
9. AB: Abelian groups and group homomorphisms.
10. LIE: Lie group and Lie group homomorphisms.
11. lie: Lie algebras and Lie algebra homomorphisms.

We use the word *class* instead of set for the collection of objects in a category is that some categories are "too large" to be considered sets. For example, in the category SET, the class of objects is the class of all sets. Any attempt to treat this class as a set in its own right leads to the Russell paradox of set theory.

Relations among morphisms are often depicted using commutative diagrams.

Definition 5.7. Morphisms can have any of the following properties. A morphism $f : X \longrightarrow Y$ is a

1. *monomorphism* (or *monic*) if $f \circ g_1 = f \circ g_2$ implies $g_1 = g_2$ for all morphisms $g_1, g_2 : Z \longrightarrow X$.
2. *epimorphism* (or *epic*) if $g_1 \circ f = g_2 \circ f$ implies $g_1 = g_2$ for all morphisms $g_1, g_2 : Y \longrightarrow Z$.
3. *bimorphism* if f is both monic and epic.
4. *isomorphism* if (defined above).
5. *endomorphism* if $X = Y$. $\text{End}(X)$ denotes the class of endomorphisms of X .
6. *automorphism* if f is both an endomorphism and an isomorphism. $\text{Aut}(X)$ denotes the class of automorphisms of X .
7. *retraction* if a right inverse of f exists, i.e. if there exists a morphism $g : Y \longrightarrow X$ such that $f \circ g = \text{Id}_Y$.

8. *section* if a left inverse of f exists, i.e. if there exists a morphism $g : Y \rightarrow X$ such that $g \circ f = \text{Id}_X$.

Theorem 5.4. *Every retraction is an epimorphism. Every section is a monomorphism. Furthermore, the following three statements are equivalent*

1. f is a monomorphism and a retraction.
2. f is an epimorphism and a section.
3. f is an isomorphism.

Definition 5.8. If \mathbf{C} and \mathbf{D} are categories, a *covariant functor* from \mathbf{C} to \mathbf{D} is a rule \mathcal{F} that assigns each object X in \mathbf{C} to an object $\mathcal{F}(X)$ in \mathbf{D} , and to each morphism $f \in \text{Hom}_{\mathbf{C}}(X, Y)$ to a morphism $\mathcal{F}(f) \in \text{Hom}_{\mathbf{D}}(\mathcal{F}(X), \mathcal{F}(Y))$, so that identities and composition are preserved.

$$\mathcal{F}(\text{Id}_X) = \text{Id}_{\mathcal{F}(X)}, \mathcal{F}(g \circ h) = \mathcal{F}(g) \circ \mathcal{F}(h)$$

There are also functors that reverse morphisms.

Definition 5.9. A *contravariant functor* \mathcal{F} from \mathbf{C} to \mathbf{D} assigns to each object X in \mathbf{C} an object $\mathcal{F}(X)$ in \mathbf{D} , and to each morphism $g \in \text{Hom}_{\mathbf{C}}(X, Y)$ a morphism $\mathcal{F}(g) \in \text{Hom}_{\mathbf{D}}(\mathcal{F}(Y), \mathcal{F}(X))$, such that

$$\mathcal{F}(\text{Id}_X) = \text{Id}_{\mathcal{F}(X)}, \mathcal{F}(g \circ h) = \mathcal{F}(g) \circ \mathcal{F}(h)$$

If the functor is understood, it is common for the morphism induced by a covariant functor to be denoted by g_* (instead of $\mathcal{F}(g)$) and that induced by a contravariant functor by g^* .

Definition 5.10. A covariant functor from any category to itself is the *identity functor*, which takes each object and each morphism to itself.

Definition 5.11. If \mathbf{C} is a category whose objects are sets with some additional structure and whose morphisms are maps preserving that structure (all categories listed in the example except for the first one), the *forgetful functor* $\mathcal{F} : \mathbf{C} \rightarrow \text{SET}$ assigns to each object its underlying set and to each morphism the same map thought of as a map between sets.

Some more interesting functors arise in the following examples.

Example 5.5. *The assignment $G \mapsto \text{Lie}(G), F \mapsto F_*$ is a covariant functor from LIE (category of Lie groups) to lie (category of Lie algebras). Note that every object G in LIE gets mapped to object $\text{Lie}(G)$ in lie, and every morphism F gets mapped to morphism F_* .*

Example 5.6. *If we define TOP_* to be the category whose objects are pointed topological spaces (topological spaces with a choice of base point in each), and whose morphisms are continuous maps taking base points to base points, then the fundamental group is a covariant functor from TOP_* to GROUP.*

Definition 5.12. The *tangent functor* is a covariant functor from the category SM of smooth manifolds to the category VB of smooth vector bundles. TO each smooth manifold M is assigns the tangent bundle $TM \rightarrow M$, and to each smooth map $F : M \rightarrow N$, it assigns the pushforward $F_* : TM \rightarrow TN$.

Definition 5.13. If F and G are covariant functors between the categories \mathbf{C} and \mathbf{D} , then a *natural transformation* η from F to G associates to every object X in \mathbf{C} a morphism $\eta_X : F(X) \rightarrow G(X)$ in \mathbf{D} such that for every morphism $f : X \rightarrow Y$ in \mathbf{C} , we have $\eta_Y \circ F(f) = G(f) \circ \eta_X$, meaning that the following diagram is commutative.

$$\begin{array}{ccc} F(X) & \xrightarrow{F(f)} & F(Y) \\ \downarrow \eta_X & & \downarrow \eta_Y \\ G(X) & \xrightarrow{G(f)} & G(Y) \end{array}$$

The two functors F and G are called *naturally isomorphic* if there exists a natural transformation from F to G such that η_X is an isomorphism for every object X in \mathbf{C} .

6 The Cotangent Bundle

Whereas tangent vectors give us a coordinate free interpretation of derivatives of curves, it turns out that derivatives of real-valued functions on a manifold are most naturally interpreted as tangent covectors. Thus, we will define the differential of a real-valued function as a covector field, similar to a coordinate-independent analogue of the classical gradient.

We will assume that the reader is familiar with the concepts of the dual space and its respective dual basis. Recall the following.

Proposition 6.1. *The dual map satisfies the following properties*

1. $(A \circ B)^* = B^* \circ A^*$
2. $(\text{Id}_V)^* : V^* \rightarrow V^*$ is the identity map of V^* .

The dual map has a nice interpretation within the context of category theory.

Corollary 6.1.1. *The assignment that sends a vector space to its dual space and a linear map to its dual map is a contravariant functor from the category of real vector spaces to itself.*

6.1 Tangent Covectors on Manifolds

Definition 6.1. Let M be a smooth manifold. For $p \in M$, we define the *cotangent space* at p , denoted by T_p^*M , to be the dual space to T_pM .

$$T_p^*M \equiv (T_pM)^*$$

Elements of T_p^*M are called *tangent covectors at p* , or just *covectors at p* . That is, if $\omega \in T_p^*M$, then

$$\omega : T_pM \rightarrow \mathbb{R}$$

If (x^i) are smooth local coordinate on an open subset $U \subset M$, then for each $p \in U$, the coordinate basis $(\partial/\partial x^i|_p)$ of T_pM induces the dual basis of T_p^*M , denoted $(\lambda^i|_p)$. Therefore, any covector $\omega \in T_p^*M$ can be written uniquely as

$$\omega = \omega_i \lambda^i|_p, \text{ where } \omega_i = \omega \left(\frac{\partial}{\partial x^i} \Big|_p \right)$$

6.1.1 Basis Transformations

Suppose that (\tilde{x}^j) is another set of smooth coordinates whose domain contains p and let $(\tilde{\lambda}^j|_p)$ denote the basis for T_p^*M dual to $(\partial/\partial \tilde{x}^j|_p)$. We can compute the components of the same covector ω with respect to the new coordinate system as follows. First, note that coordinate vector fields transform as follows:

$$\frac{\partial}{\partial x^i} \Big|_p = \frac{\partial \tilde{x}^j}{\partial x^i}(p) \frac{\partial}{\partial \tilde{x}^j} \Big|_p$$

Writing ω with respect to both bases

$$\omega = \omega_i \lambda^i|_p = \tilde{\omega}_j \tilde{\lambda}^j|_p$$

we can compute the components ω_i in terms of $\tilde{\omega}_j$.

$$\omega_i = \omega \left(\frac{\partial}{\partial x^i} \Big|_p \right) = \omega \left(\frac{\partial \tilde{x}^j}{\partial x^i}(p) \frac{\partial}{\partial \tilde{x}^j} \Big|_p \right) = \frac{\partial \tilde{x}^j}{\partial x^i}(p) \tilde{\omega}_j$$

Definition 6.2. *Covariant vectors* transform in the same way as coordinate partial derivatives. Given a vector V in coordinates with respect to basis vectors (x^i) and \tilde{V} in coordinates with respect to basis vectors (\tilde{x}^i) , we have

$$x^i = \frac{\partial \tilde{x}^j}{\partial x^i} \tilde{x}^j$$

where the indices of \tilde{x}^j cancel out with the *upper* term of the derivative meaning that if \tilde{x}^j is doubled, then the derivative is doubled and so x^i is doubled. *Contravariant vectors* transform in the opposite way.

$$\tilde{V}^j = \frac{\partial \tilde{x}^j}{\partial x^i}(p) V^i$$

where the indices of V^i cancel out with the *lower* term of the derivative, meaning that if V^i is doubled, then the derivative is halved and so \tilde{V}^j is halved.

Proposition 6.2. *It is useful to note the following:*

1. *Basis vectors of V , by definition, transform covariantly.*
2. *Coefficients of vectors in V transform contravariantly.*
3. *Basis covectors of V^* transform contravariantly.*
4. *Coefficients of covectors of V^* transform covariantly.*

Note that the use of the terms covariant and contravariant has nothing to do with the covariant and contravariant functors of category theory!

6.2 The Cotangent Bundle

Definition 6.3. Given smooth manifold M , the disjoint union

$$T^*M = \bigsqcup_{p \in M} T_p^*M$$

is called the *cotangent bundle of M* . It has a natural projection map $\pi : T^*M \rightarrow M$ sending $\omega \in T_p^*M$ to $p \in M$. In addition, given any smooth local coordinates (x^i) on $U \subset M$, for each $p \in U$ we denote the basis for T_p^*M dual to $(\partial/\partial x^i|_p)$ by $(\lambda^i|_p)$. This defines n maps $\lambda^1, \dots, \lambda^n : U \rightarrow T^*M$, called *coordinate covector fields*.

As expected, T^*M can be naturally turned into a vector bundle over M .

Proposition 6.3. *Let M be a smooth manifold and let T^*M be its cotangent bundle. With its standard projection map and the natural vector space structure on each fiber, T^*M has a unique smooth manifold structure making it into a rank- n vector bundle over M for which all coordinate covector fields are smooth local sections.*

Smooth local coordinates for M induces smooth local coordinates for its cotangent space, which in turn induces local coordinates for its cotangent bundle. That is, if (x^i) are smooth coordinates on an open set $U \subset M$, the map from $\pi^{-1}(U)$ to \mathbb{R}^{2n} given by

$$\zeta_i \lambda^i|_p \mapsto (x^1(p), \dots, x^n(p), \zeta_1, \dots, \zeta_n)$$

is a smooth coordinate chart for T^*M , called the *standard coordinates for T^*M associated with (x^i)* .

Definition 6.4. A section of T^*M is called a *covector field*, or a *differential 1-form*. The value of a covector field ω at a point $p \in M$ is denoted ω_p (since writing $w(p)$ is used to denote the action of a covector on a vector).

Definition 6.5. In any smooth local coordinates on an open set $U \subset M$, a covector field ω can be written in terms of the coordinate covector fields (λ^i) as $\omega = \omega_i \lambda^i$ for n functions $\omega_i : U \rightarrow \mathbb{R}$, called the *component functions of ω* . They are characterized by

$$\omega_i(p) = \omega_p \left(\frac{\partial}{\partial x^i} \Big|_p \right)$$

Similarly for vector fields, there are several ways to check for smoothness of a covector field.

Lemma 6.4 (Smoothness Criteria for Covector Fields). *Let M be a smooth manifold, and let $\omega : M \rightarrow T^*M$ be a rough section.*

1. *If $\omega = \omega_i \lambda^i$ is the coordinate representation for ω in any smooth chart $(U, (x^i))$ for M , then ω is smooth on U if and only if its component functions ω_i are smooth.*
2. *ω is smooth if and only if for every smooth vector field X on an open subset $U \subset M$, the function $\langle \omega, X \rangle : U \rightarrow \mathbb{R}$ defined by*

$$\langle \omega, X \rangle(p) \equiv \langle \omega_p, X_p \rangle \equiv \omega_p(X_p)$$

is smooth.

6.3 The Differential of a Function

In elementary calculus, the gradient of a smooth real valued function f on \mathbb{R}^n is defined as the vector field whose components are the partial derivatives of f . In our notation, this would read

$$\text{grad}f = \sum_{i=1}^n \frac{\partial f}{\partial x^i} \frac{\partial}{\partial x^i}$$

However, the gradient does not make coordinate independent sense. In general, although the first partial derivatives of a smooth function cannot be interpreted in a coordinate independent way as the components of a vector field, it turns out that they can be interpreted as the components of a covector field. This is the most important application of covector fields.

Definition 6.6. The covector field of f , or the *differential of f* , denoted df is defined

$$df_p(X_p) \equiv X_p f \text{ for } X_p \in T_p M$$

In general, given a vector field existing on a manifold M , applying the covector field df to it would give a scalar field on M . The definition above says that this application of the covector field df to vector field X is merely just applying the function f itself to all points in X , outputting a scalar field.

This intuition along with the following theorem leads to an alternate definition of the cotangent space.

Theorem 6.5. T_p^*M is isomorphic to $C^\infty(M)/\sim$, where \sim is the equivalence relation between curves that pass through point $p \in M$ in the same direction with the same speed. The direction of the curves in an equivalence class determines the direction of the cotangent vector and the parameterization of the curve determines its magnitude.

Definition 6.7. The vector space

$$\{(df)_p \mid f \in C^\infty(M)\}$$

is called the *cotangent space* at $p \in M$.

Lemma 6.6. The differential of a smooth function is a smooth covector field.

To see what df looks like more concretely, let us compute its coordinate representation. Let (x^i) be smooth coordinates on an open subset $U \subset M$ and let (λ^i) be the corresponding coordinates. Then, the coordinate representation of df is

$$df_p = \frac{\partial f}{\partial x^i}(p) \lambda^i \Big|_p$$

But by letting $f = x^j : U \rightarrow \mathbb{R}$, we find that

$$dx^j \Big|_p = \frac{\partial x^j}{\partial x^i} = \delta_i^j \lambda^i \Big|_p = \lambda^j \Big|_p$$

meaning that λ^j is dx^j ! Therefore, the formula for the coordinate representation of df can be rewritten as

$$df_p = \frac{\partial f}{\partial x^i}(p) dx^i \Big|_p \implies df = \frac{\partial f}{\partial x^i} dx^i$$

which is called the *total differential of f* .

Example 6.1. If $f(x, y) = x^2 y \cos x$ on \mathbb{R}^2 , then

$$\begin{aligned} df &= \frac{\partial(x^2 y \cos x)}{\partial x} dx + \frac{\partial(x^2 y \cos x)}{\partial y} dy \\ &= (2xy \cos x - x^2 y \sin x) dx + x^2 \cos x dy \end{aligned}$$

Proposition 6.7 (Properties of the Differential). *let M be a smooth manifold, and let $f, g \in C^\infty(M)$.*

1. For any $a, b \in \mathbb{R}$, we have $d(af + bg) = a df + b dg$.
2. $d(fg) = f dg + g df$
3. $d(f/g) = (g df - f dg)/g^2$ on the set where $g \neq 0$.

4. If $J \subset \mathbb{R}$ is an interval containing the image of f , and $h : J \rightarrow \mathbb{R}$ is a smooth function, then $d(h \circ f) = (h' \circ f)df$.

5. f is constant $\implies df = 0$

Clearly, if f is a smooth real-valued function on a smooth manifold M , then $df = 0$ if and only if f is constant on each component of M .

Proposition 6.8 (Derivative of a Function along a Curve). *Suppose M is a smooth manifold, $\gamma : J \rightarrow M$ is a smooth curve, and $f : M \rightarrow \mathbb{R}$ is a smooth function, then the derivative of real valued function $f \circ \gamma : \mathbb{R} \rightarrow \mathbb{R}$ is*

$$(f \circ \gamma)'(t) = df_{\gamma(t)}(\gamma'(t))$$

Proof. For any $t_0 \in J$,

$$\begin{aligned} df_{\gamma(t_0)}(\gamma'(t_0)) &= \gamma'(t_0)f \\ &= \left(\gamma_* \frac{d}{dt} \Big|_{t_0} \right) f \\ &= \frac{d}{dt} \Big|_{t_0} (f \circ \gamma) \\ &= (f \circ \gamma)'(t_0) \end{aligned}$$

■

So far, we have defined two types of derivatives for a smooth real valued function $f : M \rightarrow \mathbb{R}$ at a point $p \in M$. The first is the pushforward f_* as a linear map from $T_p M$ to $T_{f(p)}\mathbb{R}$. Later, we have defined the differential df_p as a covector at p , which is just a linear map from $T_p M$ to \mathbb{R} . But the canonical isomorphism between \mathbb{R} and its tangent space at any point leads to these two interpretations of the derivative being exactly the same.

Similarly, if γ is a smooth curve in M , we have two different meanings for the expression $(f \circ \gamma)'(t)$.

1. $f \circ \gamma$ can be interpreted as a smooth curve in \mathbb{R} , and thus $(f \circ \gamma)'(t)$ is its tangent vector at the point $(f \circ \gamma)(t) \in \mathbb{R}$, i.e. an element of the tangent space $T_{(f \circ \gamma)(t)}\mathbb{R}$.
2. Or, $f \circ \gamma$ can be considered as an ordinary function from \mathbb{R} to \mathbb{R} , with $(f \circ \gamma)'$ being just its ordinary derivative.

Either one of these two interpretations are equally correct since the derivative shown in 2 is equal to the real number $df_{\gamma(t)}(\gamma'(t))$.

6.4 Pullbacks

We know that a smooth map yields a linear map on the tangent vectors called the pushforward. Dualizing this leads to a linear map on covectors going in the opposite direction.

Definition 6.8. Let $F : M \rightarrow N$ be a smooth map, and let $p \in M$ be arbitrary. The pushforward map

$$F_* : T_p M \rightarrow T_{F(p)} N$$

yields a dual linear map

$$(F_*)^* : T_{F(p)}^* N \rightarrow T_p^* M$$

called the *pullback associated with F* . More simply, we rewrite the above as

$$F^* : T_{F(p)}^* N \rightarrow T_p^* M$$

We can see that F^* is defined by

$$(F^*\omega)(X) = \omega(F_*X) \text{ for } \omega \in T_{F(p)}^* N, X \in T_p M$$

Note that $(F^*\omega)(X)$ is a scalar field on M (since $F^*\omega$ is a covector field on M and X is a vector field on M).

We have noted that pushforwards do not always work on vector fields, but smooth covector fields always pull back to smooth covector fields. That is, given a smooth map $G : M \rightarrow N$ and a smooth covector field ω on N , we can define a covector field $G^*\omega$ on M by

$$(G^*\omega)(p) = G^*(\omega_{G(p)})$$

Notice that there is no ambiguity about what point to pull back from.

Lemma 6.9. *Let $G : M \rightarrow N$ be a smooth map, and let $f \in C^\infty(N)$ and $\omega \in \mathcal{T}^*(N)$. Then*

$$\begin{aligned} G^*df &= d(f \circ G) \\ G^*(f\omega) &= (f \circ G)G^*\omega \end{aligned}$$

Proposition 6.10. *Suppose $G : M \rightarrow N$ is smooth, and let ω be a smooth covector field on N . Then $G^*\omega$ is a smooth covector field on M .*

The formula for the pullback of a covector field with respect to smooth coordinates (x^i) on the domain and (y^j) on the range is

$$G^*\omega = G^*(\omega_j dy^j) = (\omega_j \circ G)d(y^j \circ G) = (\omega_j \circ G)dG^j$$

where G^j is the j th component function of G in these coordinates. This makes the computation of pullbacks in coordinates very simple.

Example 6.2. *Let $G : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be the map given by*

$$(u, v) = G(x, y, z) = (x^2y, y \sin z)$$

and let $\omega \in \mathcal{T}^*(\mathbb{R}^2)$ be the covector field

$$\omega = u dv + v du$$

Then, the pullback $G^*\omega$ is given by

$$\begin{aligned} G^*\omega &= (u \circ G)d(v \circ G) + (v \circ G)d(u \circ G) \\ &= (x^2y)d(y \sin z) + (y \sin z)d(x^2y) \\ &= x^2y(\sin z dy + y \cos z dz) + y \sin z(2xy dx + x^2 dy) \\ &= 2xy^2 \sin z dx + 2x^2y \sin z dy + x^2y^2 \cos z dz \end{aligned}$$

6.5 Line Integrals

We would like to make a coordinate independent sense of the line integral.

Definition 6.9. Let $[a, b] \subset \mathbb{R}$ be a compact interval, and let ω be a smooth covector field on $[a, b]$ (note that at the endpoints, smoothness of ω means that ω admits a smooth extension of to some neighborhood of $[a, b]$). Letting t denote the one standard coordinate in \mathbb{R} , then we can write ω in terms of t coordinates as

$$\omega_t \equiv f(t) dt$$

for some smooth $f : [a, b] \rightarrow \mathbb{R}$. Then, the *integral of ω over $[a, b]$* is defined

$$\int_{[a,b]} \omega \equiv \int_a^b f(t) dt$$

This definition gives rise to the following property of the line integral.

Proposition 6.11 (Diffeomorphism Invariance of the Integral). *Let ω be a smooth covector field on the compact interval $[a, b] \subset \mathbb{R}$. If $\varphi : [c, d] \rightarrow [a, b]$ is an increasing diffeomorphism (meaning that $t_1 < t_2 \implies \varphi(t_1) < \varphi(t_2)$), then*

$$\int_{[c,d]} \varphi^*\omega = \int_{[a,b]} \omega$$

Proof. Let t denote the standard coordinate on $[a, b]$ and s denote that of $[c, d]$. Then the formula for the pullback of the covector field, in coordinate expression, is

$$(\varphi^*\omega)_s = f(\varphi(s))\varphi'(s) ds$$

meaning that

$$\int_{[c,d]} \varphi^*\omega = \int_c^d f(\varphi(s))\varphi'(s) ds = \int_a^b f(t) dt = \int_{[a,b]} \omega$$

by the change of variables formula for ordinary integrals. ■

Definition 6.10. Let M be a smooth manifold. A *curve segment in M* is a continuous curve $\gamma : [a, b] \rightarrow M$ whose domain is a compact interval. It is a *smooth curve segment* if it has a smooth extension to an open interval containing $[a, b]$. A *piecewise smooth curve segment* is a curve segment $\gamma : [a, b] \rightarrow M$ with the property that there exists a finite partition of $[a, b]$ such that the image of each partition is smooth.

Now we can define line integrals over smooth covector fields on an arbitrary manifold M .

Definition 6.11. If $\gamma : [a, b] \rightarrow M$ is a smooth curve segment and ω is a smooth covector field on M , the *line integral of ω over γ* is defined to be the real number

$$\int_{\gamma} \omega \equiv \int_{[a,b]} \gamma^*\omega$$

More generally, if γ is piecewise smooth (with partitions $[a_{i-1}, a_i]$ for $i = 1, \dots, k$), then we define

$$\int_{\gamma} \omega \equiv \sum_{i=1}^k \int_{[a_{i-1}, a_i]} \gamma^*\omega$$

Note that this definition now gives rigorous meaning to classical line integrals such as

$$\int_{\gamma} P dx + Q dy$$

in \mathbb{R}^2 or

$$\int_{\gamma} P dx + Q dy + R dz$$

Proposition 6.12 (Properties of Line Integrals). *Let M be a smooth manifold. Suppose $\gamma : [a, b] \rightarrow M$ is a piecewise smooth curve segment and $\omega, \omega_1, \omega_2 \in \mathcal{T}^*(M)$.*

1. For any $c_1, c_2 \in \mathbb{R}$,

$$\int_{\gamma} (c_1\omega_1 + c_2\omega_2) = c_1 \int_{\gamma} \omega_1 + c_2 \int_{\gamma} \omega_2$$

2. If γ is a constant map, then

$$\int_{\gamma} \omega = 0$$

for all ω .

3. If $a < c < b$, then

$$\int_{\gamma} \omega = \int_{\gamma_1} \omega + \int_{\gamma_2} \omega$$

where $\gamma_1 = \gamma|_{[a,c]}$ and $\gamma_2 = \gamma|_{[c,b]}$.

The next lemma gives a useful alternative expression for the line integral of a covector field on a manifold. This is good for computations.

Lemma 6.13. *If $\gamma : [a, b] \rightarrow M$ is a piecewise smooth curve segment, the line integral of ω over γ can also be expressed as the ordinary integral*

$$\int_{\gamma} \omega = \int_a^b \omega_{\gamma(t)}(\gamma'(t)) dt$$

Example 6.3. Let $M = \mathbb{R}^2 \setminus \{0\}$, and let ω be the covector field on M given by

$$\omega \equiv \frac{x dy - y dx}{x^2 + y^2}$$

and let $\gamma : [0, 2\pi] \rightarrow M$ be the curve segment defined

$$\gamma(t) \equiv (\cos t, \sin t)$$

Since $\gamma^*\omega$ can be computed by substituting $x = \cos t$ and $y = \sin t$ everywhere in the formula for ω , we get

$$\int_{\gamma} \omega = \int_{[0, 2\pi]} \frac{\cos t(\cos t dt) - \sin t(-\sin t dt)}{\sin^2 t + \cos^2 t} = \int_0^{2\pi} dt = 2\pi$$

Another important property of line integrals is invariance under reparamaterizations.

Definition 6.12. Let $\gamma : [a, b] \rightarrow M$ and $\tilde{\gamma} : [c, d] \rightarrow M$ be smooth curve segments. If $\tilde{\gamma} = \gamma \circ \varphi$ for some diffeomorphism $\varphi : [c, d] \rightarrow [a, b]$, then it is said that $\tilde{\gamma}$ is a *reparamaterization* of γ .

If φ is an increasing function, then $\tilde{\gamma}$ is a *forward paramaterization*, and if φ is decreasing, then $\tilde{\gamma}$ is a *backward paramaterization*.

Proposition 6.14. Let M be a smooth manifold, ω a smooth covector field on M , and γ a piecewise smooth curve segment in M . For any reparamaterization $\tilde{\gamma}$ of γ , we have

$$\int_{\tilde{\gamma}} \omega = \begin{cases} \int_{\gamma} \omega, & \tilde{\gamma} \text{ is a forward paramaterization} \\ -\int_{\gamma} \omega, & \tilde{\gamma} \text{ is a backward paramaterization} \end{cases}$$

Proof. This is a direct result of the diffeomorphism invariance property of the integral. ■

There is one special case in which a line integral is trivial to compute: the line integral of a differential.

Theorem 6.15 (Fundamental Theorem of Line Integrals). Let M be a smooth manifold. Let f is a smooth real-valued function on M and let $\gamma : [a, b] \rightarrow M$ be a piecewise smooth curve segment in M . Then

$$\int_{\gamma} df = \int_{\gamma} \frac{\partial f}{\partial x^i} dx^i = f(\gamma(b)) - f(\gamma(a))$$

Proof. Suppose that γ is smooth. Then,

$$\int_{\gamma} df = \int_a^b df_{\gamma(t)}(\gamma'(t)) dt = \int_a^b (f \circ \gamma)'(t) dt = f \circ \gamma(b) - f \circ \gamma(a)$$

This naturally extends to piecewise smooth functions, since then the right hand side will be a telescoping sum in which the middle terms cancel out. ■

6.6 Conservative Covector Fields

The fundamental theorem of line integrals shows that if a covector field ω can be written as the differential of a smooth function, then the line integral can be computed extremely easily once the smooth function is known. We formally define this kind of vector field.

Definition 6.13. A smooth covector field ω on manifold M is *exact* (or an *exact differential*) on M if there exists a function $f \in C^\infty(M)$ such that $\omega = df$. The function f is called the *potential* for ω .

Note that the potential of an exact smooth covector field is not uniquely determined, but the difference between any two potentials must be constant. We now define a similar construction called the conservative covector field.

Definition 6.14. A smooth covector field ω is *conservative* if the line integral of ω over any closed piecewise smooth curve segment is 0.

It is easy to understand the following lemma given the definition of conservative covector fields.

Lemma 6.16. *A smooth covector field ω is conservative if and only if the line integral of ω depends on the endpoints of the curve. That is,*

$$\int_{\gamma} \omega = \int_{\tilde{\gamma}} \omega$$

if γ and $\tilde{\gamma}$ are piecewise smooth curve segment with the same starting and ending points.

This leads to the equivalence of exact and conservative vector fields.

Theorem 6.17. *A smooth covector field is conservative if and only if it is exact.*

7 Submersions, Immersions, and Embeddings

Because the pushforward represents the "best linear approximation" to the map near a given point, we can learn a great deal about the map itself by studying linear-algebraic properties of its pushforward at each point. The most important property is the rank of the pushforward.

7.1 Maps of Constant Rank

Definition 7.1. If $F : M \rightarrow N$ is a smooth map, the *rank of F at $p \in M$* is the rank of the linear map

$$F_* : T_p M \rightarrow T_{F(p)} M$$

In an abstract sense, it is the dimension of the image of F_* in $T_{F(p)} M$. Given any charts in the neighborhoods of p and $F(p)$, the rank of F at p is the rank of the matrix of partial derivatives of F .

Note that the rank does not depend on the choice of basis for our chart mappings.

Definition 7.2. If $F : M \rightarrow N$ has the same rank k at every point, then we can say that F has *constant rank*. That is,

$$\text{rank } F = k$$

Definition 7.3. A smooth map $F : M \rightarrow N$ is called a *submersion* if F_* is surjective at each point, or equivalently, $\text{rank } F = \dim N$.

Definition 7.4. A smooth map $F : M \rightarrow N$ is called an *immersion* if F_* is injective at each point, or equivalently, $\text{rank } F = \dim M$.

We can imagine submersions and immersions behaving locally like surjective and injective linear maps, respectively.

Definition 7.5. A *smooth embedding* is an immersion $F : M \rightarrow N$ that is also a topological embedding (i.e. a homeomorphism onto its image $F(M) \subset N$ in the subspace topology).

7.2 The Inverse Function Theorem

Definition 7.6. Let X be a metric space. A map $G : X \rightarrow X$ is said to be a *contraction* if there is a constant $\lambda < 1$ such that $d(G(x), G(y)) \leq \lambda d(x, y)$ for all $x, y \in X$. Clearly, any contraction is continuous.

Theorem 7.1 (Inverse Function Theorem on Euclidean Space). *Suppose U and V are open subsets in \mathbb{R}^n , and $F : U \rightarrow V$ is a smooth map. If $DF(p)$ is nonsingular at some point $p \in U$, then there exists connected neighborhoods $U_0 \subset U$ of p and $V_0 \subset V$ of $F(p)$ such that*

$$F|_{U_0} : U_0 \rightarrow V_0$$

is a diffeomorphism.

8 Tensors

We will always assume that the base field of every vector space mentioned is \mathbb{R} .

8.1 The Algebra of Tensors

We will assume that the reader is already familiar with the notion of tensors both as multilinear maps and as the abstract tensor product of vector spaces. There will be slight changes in notation compared to the descriptions of tensors in previous chapters.

Definition 8.1. A *covariant k -tensor* is an element of

$$T^k(V) = \bigotimes_k V^*$$

which is just a multilinear map from the Cartesian product of V 's to \mathbb{R} . A *contravariant l -tensor* is an element of

$$T_l(V) = \bigotimes_l V$$

Generally, for any nonnegative integers k, l , the space of *mixed tensors* of type (k, l) is defined as

$$T_l^k(V) \equiv \bigotimes_k V^* \otimes \bigotimes_l V$$

which is really the space of real-valued multilinear functions of k vectors and l covectors.

Strictly speaking, the $T^k(V)$ and the space of all covariant tensors are not the same, but rather naturally isomorphic to each other. However, we will treat them as the same thing.

Example 8.1 (Covariant Tensors). 1. Every linear map $\omega : V \rightarrow \mathbb{R}$ is multilinear. It is a covariant 1-tensor.

2. An inner product on V is a covariant 2-tensor, since by definition an inner product is bilinear.

3. The determinant, thought of as a function of n vectors (columns of a matrix) is a covariant n -tensor of \mathbb{R}^n .

4. Suppose $\omega, \eta \in V^*$. We define a map $\omega \otimes \eta : V \times V \rightarrow \mathbb{R}$ as

$$\omega \otimes \eta(X, Y) \equiv \omega(X) \eta(Y)$$

where the product on the right is ordinary multiplication of real numbers. The linearity of ω and η guarantees that $\omega \otimes \eta$ is a bilinear function of X and Y .

We will now provide another abstract construction of a tensor product space.

Definition 8.2. Let S be a set. A *finite formal linear combination* is a function $\mathcal{F} : S \rightarrow \mathbb{R}$ such that $\mathcal{F}(s) = 0$ for all but finitely many $s \in S$.

Definition 8.3. Let S be a set. The *free vector space on S* , denoted $\mathbb{R}\langle S \rangle$, is the set of all finite formal linear combination of elements of S with real coefficients. Under pointwise addition and scalar multiplication, $\mathbb{R}\langle S \rangle$ becomes a real vector space.

Identifying each element $x \in S$ with the function that takes the value 1 on x and 0 on all other elements of S , any element $\mathcal{F} \in \mathbb{R}\langle S \rangle$ can be written uniquely as a linear combination of these functions as the basis. This means that $\mathbb{R}\langle S \rangle$ is finite dimensional if and only if S is a finite set.

Now, we construct the tensor product space in a more abstract way.

Definition 8.4. Let V, W be finite-dimensional real vector spaces, and let \mathcal{R} be the subspace of the free vector space $\mathbb{R}\langle V \times W \rangle$ spanned by all elements of the form

$$\begin{aligned} a(v, w) - (av, w), \\ a(v, w) - (v, aw), \\ (v, w) + (v', w) - (v + v', w) \\ (v, w) + (v, w') - (v, w + w') \end{aligned}$$

for $a \in \mathbb{R}, v, v' \in V$, and $w, w' \in W$. The *tensor product* of V and W , denoted by $V \otimes W$, is the quotient space

$$V \otimes W \equiv \frac{\mathbb{R}\langle V \times W \rangle}{\mathcal{R}}$$

The equivalence class of an element (v, w) in $V \otimes W$ is denoted by $v \otimes w$.

Proposition 8.1 (Bilinearity of the Tensor Product). *The construction of the tensor product implies*

$$\begin{aligned} a(v \otimes w) &= av \otimes w = v \otimes aw \\ v \otimes w + v' \otimes w &= (v + v') \otimes w \\ v \otimes w + v \otimes w' &= v \otimes (w + w') \end{aligned}$$

Proposition 8.2 (Characteristic Property of Tensor Products). *Let V and W be finite-dimensional real vector spaces. If $A : V \times W \rightarrow X$ is a bilinear map into any vector space X , there is a unique linear map $\tilde{A} : V \otimes W \rightarrow X$ such that the following diagram commutes.*

$$\begin{array}{ccc} V \times W & \xrightarrow{A} & X \\ \downarrow \pi & \searrow \tilde{A} & \\ V \otimes W & & \end{array}$$

where $\pi(v, w) = v \otimes w$.

This is called the characteristic property because it uniquely characterizes the tensor product up to isomorphism.

Proposition 8.3 (Properties of Tensor Products). *Let V, W , and X be finite-dimensional real vector spaces.*

1. *The tensor product $V^* \otimes W^*$ is canonically isomorphic to the space $B(V, W)$ of bilinear maps from $V \otimes W$ to \mathbb{R} .*
2. *If (E_i) is a basis for V and (F_j) is a basis for W , then the set of all elements of the form $E_i \otimes F_j$ is a basis for $V \otimes W$, with $\dim V \otimes W = (\dim V)(\dim W)$.*
3. *There is a unique isomorphism $V \otimes (W \otimes X) \rightarrow (V \otimes W) \otimes X$ sending $v \otimes (w \otimes x) \mapsto (v \otimes w) \otimes x$.*

In most cases, we will concern ourselves with covariant tensors.

8.2 Tensors and Tensor Fields on Manifolds

Definition 8.5. Let M be a smooth manifold. The *bundle of covariant k -tensors on M* is defined to be

$$T^k M \equiv \bigsqcup_{p \in M} T^k(T_p M)$$

The *bundle of contravariant l -tensors* is defined

$$T_l M \equiv \bigsqcup_{p \in M} T_l(T_p M)$$

and the *bundle of mixed tensors of type (k, l)* is defined

$$T_l^k M \equiv \bigsqcup_{p \in M} T_l^k(T_p M)$$

These are all called *tensor bundles over M* .

This allows us to generalize previously defined bundles, leading to the identifications:

1. $T^0 M = T_0 M = M \times \mathbb{R}$
2. $T^1 M = T^* M$
3. $T_1 M = TM$
4. $T_0^k M = T^k M$
5. $T_l^0 M = T_l M$

Definition 8.6. A section of a tangent bundle is called a (*covariant, contravariant, or mixed*) tensor field on M . A smooth tensor field is a section that is smooth in the usual sense of smooth sections of vector bundles. The vector spaces of smooth sections of these bundles is denoted

$$\begin{aligned}\mathcal{T}^k(M) &\equiv \{\text{smooth sections of } T^k M\} \\ \mathcal{T}_l(M) &\equiv \{\text{smooth sections of } T^l M\} \\ \mathcal{T}_l^k(M) &\equiv \{\text{smooth sections of } T_l^k M\}\end{aligned}$$

In any smooth local coordinates (x^i) , sections of these bundles can be written (using the summation convention) as

$$\sigma = \begin{cases} \sigma_{i_1 \dots i_k} dx^{i_1} \otimes \dots \otimes dx^{i_k}, & \sigma \in \mathcal{T}^k(M) \\ \sigma^{j_1 \dots j_l} \frac{\partial}{\partial x^{j_1}} \otimes \dots \otimes \frac{\partial}{\partial x^{j_l}}, & \sigma \in \mathcal{T}_l(M) \\ \sigma_{i_1 \dots i_k}^{j_1 \dots j_l} dx^{i_1} \otimes \dots \otimes dx^{i_k} \otimes \frac{\partial}{\partial x^{j_1}} \otimes \dots \otimes \frac{\partial}{\partial x^{j_l}}, & \sigma \in \mathcal{T}_l^k(M) \end{cases}$$

The functions $\sigma_{i_1 \dots i_k}^{j_1 \dots j_l} : M \rightarrow \mathbb{R}$ are called the *component functions* of σ .

Lemma 8.4. Let M be a smooth vector field. The following are equivalent.

1. σ is smooth.
2. In any smooth coordinate chart, the component functions of σ are smooth.
3. If X_1, \dots, X_k are smooth vector fields defined on any open subset $U \subset M$, then the function $\sigma(X_1, \dots, X_k) : U \rightarrow \mathbb{R}$ defined by

$$\sigma(X_1, \dots, X_k)(p) \equiv \sigma_p(X_1|_p, \dots, X_k|_p)$$

is smooth.

8.2.1 Pullbacks

8.3 Symmetric Tensors

Definition 8.7. Let V be a finite-dimensional vector space. A covariant k -tensor T on V is said to be *symmetric* if its value is unchanged by interchanging any pair of arguments.

$$T(X_1, \dots, X_i, \dots, X_j, \dots, X_k) = T(X_1, \dots, X_j, \dots, X_i, \dots, X_k)$$

Since the set of transpositions form the generating set of the set of all permutations of n elements, a symmetric tensor is invariant under any permutation of its arguments.

Definition 8.8. The set of symmetric covariant k -tensors on V is denoted $\Sigma^k(V)$, which is a vector subspace of $T^k(V)$. There is a natural surjective projection

$$\text{Sym} : T^k(V) \rightarrow \Sigma^k(V)$$

called *symmetrization*. Given a k -tensor T and a permutation $\sigma \in S_k$ (S_k is the symmetric group of k elements), we define the symmetrization of tensor T by

$$\text{Sym}(T) \equiv \frac{1}{k!} \sum_{\sigma \in S_k} T(X_{\sigma(1)}, \dots, X_{\sigma(k)})$$

Note that if S and T are symmetric tensors on V , then $S \otimes T$ is not symmetric in general. But we can simply define a new product that does produce a symmetric product.

Definition 8.9. Given $S \in \Sigma^k(V)$ and $T \in \Sigma^l(V)$, their *symmetric product* is the $(k+l)$ -tensor ST defined

$$ST \equiv \text{Sym}(S \otimes T)$$

We can explicitly define this as

$$ST(X_1, \dots, X_{k+l}) \equiv \frac{1}{(k+l)!} \sum_{\sigma \in S_{k+l}} S(X_{\sigma(1)}, \dots, X_{\sigma(k)}) T(X_{\sigma(k+1)}, \dots, X_{\sigma(k+l)})$$

Note that the symmetric product (written using juxtaposition of tensors above) is the same product as the product " \odot " mentioned in the linear algebra chapter.

Proposition 8.5 (Properties of the Symmetric Product). *We list two additional important properties.*

1. *The symmetric product is symmetric and bilinear. For all symmetric tensors R, S, T and all $a, b \in \mathbb{R}$,*

$$ST = TS$$

$$(aR + bS)T = aRT + bST = T(aR + bS)$$

2. *If ω and η are covectors, then*

$$\omega\eta = \frac{1}{2}(\omega \otimes \eta + \eta \otimes \omega)$$

Definition 8.10. A *symmetric tensor field* on a manifold is a covariant tensor field whose value at any point is a symmetric tensor.

8.4 Riemannian Metrics

Definition 8.11. A *Riemannian metric* on manifold M is a smooth symmetric 2-tensor field that is positive definite at each point. A *Riemannian manifold* is a pair (M, g) where M is a smooth manifold and g is a Riemannian metric on M .

Note that a Riemannian metric is not the same as a metric on a vector space.

Proposition 8.6 (Existence of Riemannian Metrics). *Every smooth manifold admits a Riemannian metric.*

If g is a Riemannian metric on M , then for each $p \in M$, g_p is an inner product on T_pM . For $X, Y \in T_pM$, the expression $\langle X, Y \rangle_g$ is used to denote the real number $g_p(X, Y)$. Furthermore, in any smooth local coordinates (x^i) , a Riemannian metric can be written

$$g = g_{ij}dx^i \otimes dx^j$$

where g_{ij} is a symmetric positive definite matrix of smooth functions. The symmetry of g also allows us to write g as

$$\begin{aligned} g &= g_{ij}dx^i \otimes dx^j \\ &= \frac{1}{2}(g_{ij}dx^i \otimes dx^j + g_{ji}dx^i \otimes dx^j) \\ &= \frac{1}{2}(g_{ij}dx^i \otimes dx^j + g_{ij}dx^i \otimes dx^j) \\ &= g_{ij}dx^i dx^j \end{aligned}$$

Example 8.2. *The simplest example of a Riemannian metric is the Euclidean metric \bar{g} on \mathbb{R}^n , defined in standard coordinates as*

$$\bar{g}(x, y) \equiv \delta_{ij}dx^i dx^j$$

where δ_{ij} is the Kronecker delta. It is common to use the notation ω^2 for the symmetric product of tensor ω with itself, so the Euclidean metric can also be written

$$\bar{g} = (dx^1)^2 + \dots + (dx^n)^2$$

8.4.1 Pseudo-Riemannian Metrics

Relaxing the requirement that the metric be positive definite results in a generalization of the Riemannian metric.

Definition 8.12. A 2-tensor g on a vector space V is said to be nondegenerate if $g(X, Y) = 0$ for all $Y \in V$ if and only if $X = 0$.

Just as any inner product can be transformed to the Euclidean one by switching to an orthonormal basis, every nondegenerate symmetric 2-tensor can be transformed by a change of basis to one with a diagonal matrix of diagonal entries ± 1 . However, the number of -1 's and $+1$'s are invariant under a choice of basis. That is, the *signature* of g is an invariant of g .

9 Differential Forms

We have introduced line integrals of covector fields, which generalized ordinary integrals to curves in manifolds. Now, we will generalize the theory of multiple integrals over manifolds.

9.1 The Algebra of Alternating Tensors

Definition 9.1. A covariant k -tensor T on a finite-dimensional vector space V is said to be *alternating* if it has the property

$$T(X_1, \dots, X_i, \dots, X_j, \dots, X_k) = -T(X_1, \dots, X_j, \dots, X_i, \dots, X_k)$$

An alternating k -tensor is also called a *k-covector*. Note that this

Bilinearity and the alternating properties of the alternating k -tensor indicates that it is a good measure of the *signed volume* of a parallelepiped. The following lemma also supports this notion, since a m -dimensional parallelepiped in an n -dimensional space has volume 0 when $m < n$.

Lemma 9.1. Suppose Ω is a k -tensor on a vector space V with the property that $\Omega(X_1, \dots, X_k) = 0$ whenever X_1, \dots, X_k is linearly dependent. Then Ω is alternating.

Proof. By bilinearity, the hypothesis says that Ω gives the value 0 whenever two arguments are the same. So,

$$\begin{aligned} 0 &= \Omega(X_1, \dots, X_{i+j}, \dots, X_{i+j}, \dots, X_n) \\ &= \Omega(X_1, \dots, X_i, \dots, X_i, \dots, X_n) + \Omega(X_1, \dots, X_i, \dots, X_j, \dots, X_n) \\ &\quad + \Omega(X_1, \dots, X_j, \dots, X_i, \dots, X_n) + \Omega(X_1, \dots, X_j, \dots, X_j, \dots, X_n) \\ &= \Omega(X_1, \dots, X_i, \dots, X_{i+j}, \dots, X_n) + \Omega(X_1, \dots, X_j, \dots, X_{i+j}, \dots, X_n) \end{aligned}$$

which means that

$$\Omega(X_1, \dots, X_i, \dots, X_{i+j}, \dots, X_n) = -\Omega(X_1, \dots, X_j, \dots, X_{i+j}, \dots, X_n)$$

■

Because of these properties, alternating tensor fields are good candidates for objects that can be integrated in a coordinate-independent way.

Proposition 9.2. The following are equivalent for a covariant k -tensor T .

1. T is alternating.
2. For any vectors X_1, \dots, X_n ,

$$T(X_{\sigma(1)}, \dots, T(X_{\sigma(n)})) = (\text{sgn}(\sigma)T(X_1, \dots, X_n))$$
3. T gives zero whenever two of its arguments are equal.
4. $T = 0$ whenever its arguments are linearly dependent.

Note that any 2-tensor can be expressed as a sum of its symmetric components and alternating components, since

$$\begin{aligned} T(X, Y) &= \frac{1}{2}(T(X, Y) - T(X, Y)) + \frac{1}{2}(T(X, Y) + T(Y, X)) \\ &= A(X, Y) + S(X, Y) \end{aligned}$$

where $A(X, Y) = \frac{1}{2}(T(X, Y) - T(X, Y))$ is alternating and $S(X, Y) = \frac{1}{2}(T(X, Y) + T(X, Y))$ is symmetric. However, this is not true for tensors of higher rank.

Note that S is just defined as $\text{Sym } T$, the symmetric projection of T . We can define a similar projection

$$\text{Alt} : T^k(V) \longrightarrow \Lambda^k(V)$$

called the *alternating projection*, defined

$$\text{Alt } T \equiv \frac{1}{k!} \sum_{\sigma \in S_k} (\text{sgn } \sigma)(X_{\sigma(1)}, \dots, X_{\sigma(k)})$$

Example 9.1. Let T be a 1-tensor. Then $\text{Alt } T = T$. If T is a 2-tensor, then

$$\text{Alt } T(X, Y) = \frac{1}{2}(T(X, Y) - T(Y, X))$$

If T is a 3-tensor,

$$\begin{aligned} \text{Alt } T(X, Y, Z) &= \frac{1}{6}(T(X, Y, Z) + T(Y, Z, X) + T(Z, X, Y) \\ &\quad - T(Y, X, Z) - T(X, Z, Y) - T(Z, Y, X)) \end{aligned}$$

Lemma 9.3 (Properties of the Alternating Projection). For any tensor T , $\text{Alt } T$ is alternating. T is alternating if and only if $\text{Alt } T = T$.

Proposition 9.4. Let V be an n -dimensional vector space. If (ε^i) is any basis for V^* , then for each positive integer $k \leq n$, the collection of k -covectors

$$\mathcal{E} = \{\varepsilon^I \mid I \text{ is an increasing multi-index of length } k\}$$

is a basis for $\Lambda^k(V)$. Therefore,

$$\dim \Lambda^k(V) = \binom{n}{k}$$

and if $k > n$, then $\dim \Lambda^k(V) = 0$.

9.2 The Wedge Product

Definition 9.2. If $\omega \in \Lambda^k(V)$ and $\eta \in \Lambda^l(V)$, the *wedge product*, or *exterior product* of ω and η is the alternating $(k+l)$ -tensor

$$\omega \wedge \eta \equiv \frac{(k+l)!}{k!l!} \text{Alt}(\omega \otimes \eta)$$

Lemma 9.5. Let $(\varepsilon^1, \dots, \varepsilon^n)$ be a basis for V^* . For any multi-indices $I = (i_1, \dots, i_k)$ and $J = (j_1, \dots, j_l)$,

$$\varepsilon^I \wedge \varepsilon^J = \varepsilon^{IJ}$$

where IJ is the multi-index $(i_1, \dots, i_k, j_1, \dots, j_l)$ obtained by concatenating I and J .

Proposition 9.6 (Properties of the Wedge Product). We list properties of the wedge product.

1. *Bilinearity.*

$$\begin{aligned} (a\omega + a'\omega') \wedge \eta &= a(\omega \wedge \eta) + a'(\omega' \wedge \eta) \\ \eta \wedge (a\omega + a'\omega') &= a(\eta \wedge \omega) + a'(\eta \wedge \omega') \end{aligned}$$

2. *Associativity.*

$$\omega \wedge (\eta \wedge \zeta) = (\omega \wedge \eta) \wedge \zeta$$

3. *Anticommutativity.*

$$\omega \wedge \eta = (-1)^{kl} \eta \wedge \omega$$

4. If $(\varepsilon^1, \dots, \varepsilon^n)$ is any basis for V^* and $I = (i_1, \dots, i_k)$ is any multi-index, then

$$\varepsilon^{i+1} \wedge \dots \wedge \varepsilon^{i_k} = \varepsilon^I$$

5. For any covectors $\omega^1, \dots, \omega^k$ and vectors X_1, \dots, X_k ,

$$\omega^1 \wedge \dots \wedge \omega^k(X_1, \dots, X_k) = \det(\omega^j(X_i))$$

Definition 9.3. For any n -dimensional vector space V , the vector space $\Lambda^*(V)$ is defined

$$\Lambda^*(V) = \bigoplus_{k=0}^n \Lambda^k(V)$$

called the *exterior algebra* of V .

Definition 9.4. An algebra A is said to be *graded* if it has a direct sum decomposition $A = \bigoplus_k A^k$ such that the product satisfies $(A^k)(A^l) \subset A^{k+l}$.

A graded algebra is *anticommutative* if the product satisfies $ab = (-1)^{kl}ba$ for $a \in A^k, b \in A^l$.

Clearly, $\Lambda^*(V)$ is an anticommutative graded algebra.

9.3 Differential Forms on Manifolds

Definition 9.5. Given an n -dimensional smooth manifold M , the subset of $T^k M$ consisting of alternating tensors is defined

$$\Lambda^k M = \bigsqcup_{p \in M} \Lambda^k(T_p M)$$

It is a smooth subbundle of $T^k M$.

Definition 9.6. A section of $\Lambda^k M$ is called a *differential k -form*, or just a *k -form*. In other words, it is a continuous tensor field whose value at each point is an alternating tensor. The integer k is called the *degree* of the form.

Definition 9.7. The vector space of smooth sections of $\Lambda^k M$ is denoted $\mathcal{A}^k M$.

In any smooth chart, a k -form ω can be written locally as

$$\omega = \sum_I \omega_I dx^{i_1} \wedge \dots \wedge dx^{i_k} = \sum_I \omega_I dx^I$$

where the coefficients ω_I are continuous functions defined on the coordinate domain. We use dx^I as an abbreviation for $dx^{i_1} \wedge \dots \wedge dx^{i_k}$. Interpreting the basis forms as forms themselves,

$$dx^{i_1} \wedge \dots \wedge dx^{i_k} \left(\frac{\partial}{\partial x^{j_1}}, \dots, \frac{\partial}{\partial x^{j_k}} \right) = \delta_J^I$$

Thus, the component functions are determined by

$$\omega^I = \omega \left(\frac{\partial}{\partial x^{i_1}}, \dots, \frac{\partial}{\partial x^{i_k}} \right)$$

Example 9.2. On \mathbb{R}^3 , some examples of smooth 2-forms are given by

$$\begin{aligned} \omega &= (\sin xy) dy \wedge dz \\ \eta &= dx \wedge dy + dx \wedge dz + dy \wedge dz \end{aligned}$$

A 0-form is just a continuous real-valued function, and a 1-form is a covector field. Every n -form \mathbb{R}^n is a continuous real-valued function times $dx^1 \wedge \dots \wedge dx^n$.

We can take the vector spaces of smooth sections of $\Lambda^k(M)$ of all the degrees k and direct sum them to create an algebra.

Definition 9.8. Defining the wedge product of two differential forms pointwise

$$(\omega \wedge \eta)_p \equiv \omega_p \wedge \eta_p$$

we can see that the wedge product of a k -form with an l -form is a $(k+l)$ -form. Defining

$$\mathcal{A}^*(M) \equiv \bigoplus_{k=0}^n \mathcal{A}^k(M)$$

equipped with the wedge product turns this set into a associative, anticommutative graded algebra.

If $F : M \rightarrow N$ is a smooth map and ω is a smooth differential form on N , the pullback $F^*\omega$ is a smooth differential form on M , defined as for any smooth tensor field

$$(F^*\omega)_p(X_1, \dots, X_k) \equiv \omega_{F(p)}(F_*X_1, \dots, F_*X_k)$$

If $i : N \rightarrow M$ is the inclusion map of an immersed submanifold, then we denote it as $\omega|_N$ for $i^*\omega$.

The following lemma gives a computational rule for pullbacks of differential forms. It can also be used to compute the expression for a differential form in another smooth chart.

Lemma 9.7. Suppose $F : M \rightarrow N$ is smooth.

1. $F^* : \mathcal{A}^k(N) \rightarrow \mathcal{A}^k(M)$ is linear.

$$2. F^*(\omega \wedge \eta) = (F^*\omega) \wedge (F^*\eta)$$

3. In any smooth chart

$$F^*\left(\sum_I \omega_I dy^{i_1} \wedge \dots \wedge dy^{i_k}\right) = \sum_I (\omega_I \circ F) d(y^{i_1} \circ F) \wedge \dots \wedge d(y^{i_k} \circ F)$$

Proposition 9.8. Let $F : M \rightarrow N$ be a smooth map between n -manifolds. If (x^i) and (y^j) are smooth coordinates on open sets $U \subset M$ and $V \subset N$, respectively, and u is a smooth real-valued function on V , then the following holds on $U \cap F^{-1}(V)$.

$$F^*(udy^1 \wedge \dots \wedge dy^n) = (u \circ F)(\det DF) dx^1 \wedge \dots \wedge dx^n$$

where DF represents the matrix of partial derivatives of F in coordinates.

Corollary 9.8.1. If $(U, (x^i))$ and $(\tilde{U}, (\tilde{x}^j))$ are overlapping smooth coordinate charts on M , then the following identity holds on $U \cap \tilde{U}$:

$$d\tilde{x}^1 \wedge \dots \wedge d\tilde{x}^n = \det\left(\frac{\partial \tilde{x}^j}{\partial x^i}\right) dx^1 \wedge \dots \wedge dx^n$$

9.4 Exterior Derivatives

Theorem 9.9 (The Exterior Derivative). For every smooth manifold M , there are unique linear maps

$$d : \mathcal{A}^k(M) \rightarrow \mathcal{A}^{k+1}(M)$$

defined for each integer $k \geq 0$ and satisfying the following conditions:

1. If f is a smooth real-valued function (i.e. a 0-form), then df is the differential of f , defined as

$$df(X) \equiv Xf$$

2. If $\omega \in \mathcal{A}^k(M)$ and $\eta \in \mathcal{A}^l(M)$, then

$$d(\omega \wedge \eta) = d\omega \wedge \eta + (-1)^k \omega \wedge d\eta$$

3. $d \circ d = 0$

The operator d also satisfies the following properties:

1. In every smooth coordinate chart, d is given by

$$d\left(\sum_J \omega_J dx^J\right) = \sum_J d\omega_J \wedge dx^J$$

2. d is local. That is, if $\omega = \omega'$ on an open set $U \subset M$, then $d\omega = d\omega'$ on U .

3. d commutes with restriction. That is, if $U \subset M$ is any open set, then

$$d(\omega|_U) = (d\omega)|_U$$

Lemma 9.10 (Naturality of the Exterior Derivative). If $G : M \rightarrow N$ is a smooth map, then the pullback map

$$G^* : \mathcal{A}^k(N) \rightarrow \mathcal{A}^k(M)$$

commutes with d . That is, for all $\omega \in \mathcal{A}^k(N)$,

$$G^*(d\omega) = d(G^*\omega)$$

10 Orientations

10.1 Orientations on Vector Spaces

In order to properly define the integration of k -forms in a way that is consistent with signed volume of a parallelepiped, we must properly define the orientation of certain vector spaces.

We could try this using a certain choice of basis. For example, given a choice of *ordered* basis vectors $\{f_1, \dots, f_n\}$, we can define the orientation of the vector space spanned by these vectors by computing the sign of the determinant of the matrix with column vectors f_i . However, since abstract vector spaces have no canonical vector spaces, we cannot say which vector spaces have the "positive orientation" or is "right-handed." However, we can *compare* whether two bases have a consistent orientation. Thus, we are led to the following definition.

Definition 10.1. Let V be a vector space of dimension $n \geq 1$. We say that two ordered bases

$$\begin{aligned} &(E_1, \dots, E_n) \\ &(\tilde{E}_1, \dots, \tilde{E}_n) \end{aligned}$$

are *consistently oriented* if the transition matrix (B_i^j) , defined by the equation

$$E_i = B_i^j \tilde{E}_j$$

has positive determinant.

Definition 10.2. Given vector space V with $\dim V \geq 1$, the *orientation* of V is an equivalence class of ordered bases. If (E_1, \dots, E_n) is any ordered basis for V , we denote that orientation that it determines by (the equivalence class)

$$[E_1, \dots, E_n]$$

A vector space together with a choice is called an *oriented vector space*. If V is oriented, then any ordered basis (E_1, \dots, E_n) that is in the given orientation is said to be *oriented* or *positively oriented*. Any basis that is not in the given orientation is said to be *negatively oriented*.

For the special case of a 0-dimensional vector space V , we define an orientation of V to be simply a choice of one of the number ± 1 .

Example 10.1. The orientation $[e_1, \dots, e_n]$ of \mathbb{R}^n determined by the standard basis is called the *standard orientation*. The standard orientation for

1. \mathbb{R} is the unit vector pointing to the right.
2. \mathbb{R}^2 is one which the rotation from the first vector to the second is counterclockwise.
3. \mathbb{R}^3 is one that has a right-handed orientation.

There is an important connection between orientation and alternating tensors, expressed in the following lemma.

Lemma 10.1. Let V be a vector space of dimension ≥ 1 , and let Ω is a nonzero element of $\Lambda^n(V)$. The set of ordered bases (E_1, \dots, E_n) such that $\Omega(E_1, \dots, E_n) > 0$ is an orientation of V .

Definition 10.3. if v is an oriented vector space and Ω is an n -covector that determines the orientation of V as described in the previous lemma, it is said that Ω is an *oriented* (or *positively oriented*) n -covector.

Example 10.2. The n -covector $e^1 \wedge \dots \wedge e^n$ is *positively oriented* for the standard orientation on \mathbb{R}^n .

10.2 Orientations on Manifolds

Definition 10.4. Let M be a smooth manifold with a given pointwise orientation. It is said that a local frame (E_i) for M is (*positively*) *oriented* if $(E_1|_p, \dots, E_n|_p)$ is a positively oriented basis for $T_p M$ at each point $p \in M$. A *negatively oriented manifold* is defined analogously.

Definition 10.5. A pointwise orientation is said to be *continuous* if every point of M is in the domain of an oriented local frame. An *orientation* of M is a continuous pointwise orientation. An *oriented manifold* is a smooth manifold together with a choice of orientation. It is said that M is *orientable* if there exists an orientation for it, and *nonorientable* if there isn't.